



Quantifying the fatal and non-fatal burden of disease associated with child growth failure, 2000–2023: a systematic analysis from the Global Burden of Disease Study 2023



GBD 2023 Child Growth Failure Collaborators*

Lancet Child Adolesc Health
2026; 10: 22–38

*Collaborators listed at the end of the Article

Correspondence to:
Professor Robert C Reiner Jr,
Institute for Health Metrics and
Evaluation, University of
Washington, Seattle, WA 98105,
USA
bcreiner@uw.edu

Summary

Background Child growth failure (CGF), which includes underweight, wasting, and stunting, is among the factors most strongly associated with mortality and morbidity in children younger than 5 years worldwide. Poor height and bodyweight gain arise from a variety of biological and sociodemographic factors and are associated with increased vulnerability to infectious diseases. We used data from the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) 2023 to estimate CGF prevalence, the risk of infectious diseases associated with CGF, and the disease mortality, morbidity, and overall burden associated with CGF.

Methods In this analysis we estimated the all-cause and cause-specific (diarrhoea, lower respiratory tract infections, malaria, and measles) disability-adjusted life-years (DALYs) lost and mortality associated with stunting, wasting, underweight, and CGF in aggregate. We combined the burden associated with mild, moderate, and severe forms of CGF: stunting was defined as height-for-age Z scores (HAZ) less than -1 , underweight was defined as weight-for-age Z scores (WAZ) less than -1 , and wasting was defined as weight-for-height Z scores (WHZ) less than -1 , according to WHO Child Growth Standards. Population-level continuous distributions of HAZ, WAZ, and WHZ were estimated for 2000 to 2023 using data from surveys, literature, and individual-level study data. The risk of incidence of, and mortality due to, diarrhoea, lower respiratory infections, malaria, and measles was separately estimated in a meta-regression framework from longitudinal cohort data for Z scores less than -1 . Finally, fatal outcomes associated with these diseases were estimated with vital registration, verbal autopsy, and case-fatality data, while non-fatal outcomes were estimated with surveys as well as health-care utilisation and case reporting data. The exposure prevalence and relative risk estimates were from continuous distributions, allowing for direct assessment of the attributable fractions for mild, moderate, and severe stunting, underweight, wasting, and the combined impact of child growth failure within populations. All estimates were age-specific, sex-specific, geography-specific, and year-specific.

Findings We estimated that, in children younger than 5 years in 2023, CGF was associated with 79.4 million (95% uncertainty interval [UI] 47.0–106) DALYs lost and 880 000 (517 000–1 170 000) deaths. This represented 17.9% (10.6–23.8) of 444 million (434–457) total under-5 DALYs and 18.8% (11.1–25.0) of all 4.67 million (4.59–4.75) under-5 deaths. Compared to stunting (33.0 million [24.1–42.2] DALYs, 373 000 [272 000–477 000] deaths) and wasting (39.2 million [23.8–53.0] DALYs, 428 000 [256 000–583 000] deaths), childhood underweight was associated with the largest share of CGF-related disease burden: 52.2 million (21.9–75.1) DALYs and 573 000 (236 000–824 000) deaths in children younger than 5 years in 2023.

Interpretation CGF remains a leading factor associated with death and disability in children younger than 5 years, despite global attention and focused interventions to reduce the prevalence of associated CGF indicators. Our findings underscore the need for policies, strategies, and interventions that focus on all indicators of CGF to reduce its associated health burden.

Funding Gates Foundation.

Copyright © 2025 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Child growth failure (CGF), characterised by poor linear height and bodyweight gain, represents a complex interplay of nutritional, maternal, developmental, socio-economic, environmental, and health-care factors.¹ Children who experience growth failure have a greater

infectious disease burden, including risk of mortality, than those who do not.^{2–5} CGF is associated with lifelong consequences such as cognitive and metabolic impairment, which might contribute to poorer educational performance and lost opportunities.^{1,6} Therefore, quantifying its prevalence and associated

Research in context

Evidence before this study

Even with decades of substantial global investment, child growth failure (CGF) remains a leading risk factor associated with death and disability in children younger than 5 years. In the most recent global estimates of CGF burden, the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) 2021 estimated that all CGF (mild, moderate, and severe) was associated with 774 000 deaths and 70.2 million DALYs in children younger than 5 years in 2021. Despite the crucial importance of understanding the causes and relative risks of negative health outcomes associated with CGF, global estimates of the increased relative risks of negative health outcomes associated with CGF typically rely on a synthetic review of 11 studies that were conducted between 1977 and 1997.

Added value of this study

Using updated estimates from GBD 2023 of both the relative risks of CGF associated with health outcomes as well as the proportion of children experiencing CGF, we present estimates of the global burden associated with CGF aggregated across mild, moderate, and severe forms for each of three CGF indicators: stunting (low height-for-age), underweight (low

weight-for-age), and wasting (low weight-for-height). For each indicator, our estimates represent burden estimates for children that are at least 1 SD below the WHO Child Growth Standards. The length and quality of the studies included in the re-estimation of relative risks allowed for distinct quantification of the morbidity and mortality risk associated with each CGF indicator for lower respiratory infections, diarrhoea, measles, and malaria. In particular, using relative risks based on increased mortality alone, we found that more than 18% of all under-5 deaths were associated with CGF, with almost as many deaths associated with stunting as with wasting.

Implications of all the available evidence

Previous estimates of CGF identified wasting as the primary CGF risk factor associated with disease burden. We found that stunting is associated with almost as much of the global CGF burden as wasting and that stunting prevalence has decreased at a far slower rate than wasting over the past decades. Given the previously observed difficulty in a child recovering from stunting after their third month of life, our estimates should serve as a call to refocus efforts to avert CGF at the earliest stages of onset.

disease burden on a global scale is of crucial importance for devising targeted interventions and policies aimed at alleviating its associated burden and generating advocacy for funding prevention and treatment.

Since 1990, substantial strides have been made in reducing childhood mortality and improving overall child growth and health,⁷ despite some setbacks caused by disruptions in health-care services and treatment of acute malnutrition due to the COVID-19 pandemic.⁸ However, many countries are not on track to meet the UN Sustainable Development Goals for reducing child mortality and the prevalence of CGF, and there is substantial overlap in the countries that have high child mortality and CGF.^{7,9} In 2019, more than half of all under-5 deaths globally were attributable to malnutrition, including CGF, low birthweight and short gestation, suboptimal breastfeeding, vitamin A deficiency, and zinc deficiency.⁷ This is in part because growth failure is a leading risk factor associated with primary infectious causes of death among children younger than 5 years.³

Numerous interventions exist that can prevent CGF, such as increased rates of exclusive breastfeeding¹⁰ and improved micronutrient uptake,¹¹ but the effectiveness of these interventions varies across CGF indicators¹¹ as well as by the age of the child.¹² As outlined in the most recent WHO guidelines, severe acute malnutrition in children from 0 to 59 months of age can frequently be managed by community health workers and increased monitoring of the child's health.¹³ Although community case management can also help reduce stunting, the likelihood of successfully reversing stunting decreases

as the infant grows older,¹⁴ indicating that identifying children whose growth is faltering as early as possible is critical. Substantial evidence suggests that short gestation, low birthweight, and factors influencing a fetus's development in utero drive many growth outcomes in the early years of life and that the ideal time to intervene on growth failure might be before the child is born¹⁵ or even conceived.¹⁶ Finally, although CGF is a risk factor associated with more severe outcomes for various infectious diseases, enteric infections can also alter intestinal absorption rates, increasing the likelihood of CGF and creating a so-called vicious cycle.¹⁷ Improved integrated case management and immediate access to treatments such as oral rehydration solution can reduce the severity of disease and in turn decrease the likelihood of growth failure.¹⁷ Proximal and effective interventions with strong evidence of impact are described in appendix 1 (table S7). A concerted effort by the global community to increase detection of CGF and access to treatment has led to substantial declines in CGF over the past few decades.

Many existing estimates of CGF focus primarily on the prevalence of moderate and severe stunting, underweight, and wasting, defined by the proportions of children less than two Z scores from the median age-specific and sex-specific global growth standards for height-for-age (HAZ) for stunting, weight-for-age (WAZ) for underweight, and weight-for-height (WHZ) for wasting.^{4,18} Measuring the prevalence of these indicators is convenient for comparing populations between geographical regions and over time and for defining and

See Online for appendix 1

Panel: Rationale for modelling all child growth failure (mild, moderate, and severe)

We modelled all child growth failure (mild, moderate, and severe) at height-for-age Z scores (HAZ), weight-for-age Z scores (WAZ), and weight-for-height Z scores (WHZ) less than -1 to capture a broader distribution of risk than the conventional Z score less than -2 threshold. Evidence suggests that the risks of infectious morbidity and mortality increase even with mild growth deficits (Z score <-1), and continuous risk curves allow more accurate estimation of the attributable burden. This approach aligns with Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) methodology, which emphasises modelling the full distribution of exposure rather than categorical cut-offs.

For the GBD 2023 Sources Tool see <https://ghdx.healthdata.org/gbd-2023/sources>

monitoring progress towards global goals.¹⁹ However, focusing exclusively on these prevalence estimates might obscure meaningful changes in population distributions and does not connect CGF with disease burden and mortality.²⁰

This analysis extends on previous estimates of the prevalence of CGF by including all forms of CGF (mild, moderate, and severe; panel), quantifying infectious-disease-specific attributable fractions from previously estimated continuous distributions of growth indicators and newly estimated continuous relative risks for incidence and mortality due to diarrhoea, lower respiratory infections, measles, and malaria.^{4,18,20} In this study, based on the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) 2023, we present comprehensive estimates of disability-adjusted life-years (DALYs) and deaths associated with CGF at the global, regional, and national level, including changes over time. Additionally, we describe updated cause-specific risks of incidence and mortality associated with each CGF indicator and the associated attributable burden. The resulting estimates of the disease burden associated with CGF are available for 204 countries and territories, five age groups younger than 5 years, by sex, from 1990 to 2023. This manuscript was produced as part of the GBD Collaborator Network and in accordance with the GBD Protocol.²¹

Methods

Overview

We used four main methodological steps to estimate the burden of all CGF (mild, moderate, and severe). To estimate the attributable fractions for each of the three risk factor components of CGF modelled in GBD—stunting, underweight, and wasting—for diarrhoea, lower respiratory infections, malaria, and measles, we produced estimates of exposure and estimates of the risk of disease given varying levels of exposure. Finally, we multiplied those attributable fractions by cause-specific cases, deaths, and DALYs to produce our results for CGF burden. Each step is described briefly here with specific

references for further information. The analyses in this study follow the methods published by the GBD 2023 Disease and Injury and Risk Factor Collaborators.²² They have been described in full elsewhere²² and are summarised below.

Estimation of the population-level distributions of HAZ, WAZ, and WHZ

A detailed description of the methods used to estimate continuous distributions for each of the CGF indicators is provided in the study by Fitzgerald and colleagues.²⁰ We included data from various sources, such as population-representative surveys, administrative data, and published scientific literature. More than 1700 sources were used in this model; citations and metadata for all sources used in the analysis are available for download from the GBD 2023 Sources Tool. A spatiotemporal Gaussian process regression (ST-GPR) model, a methodology used across many models in GBD that leverages evidence across time and space to produce estimates for each indicator, age group, sex, year, and location, was used to make predictions of the mean HAZ, WAZ, and WHZ and the prevalence of moderate (<-2 Z scores) and severe (<-3 Z scores) growth failure by age, sex, year, and location. This model includes linear effects on maternal care and immunisation, Healthcare Access and Quality (HAQ) Index, prevalence of severe anaemia, the Socio-demographic Index (SDI), age-specific and sex-specific unsafe sanitation summary exposure value, and all-age energy (kilocalories per person per day) as covariates.

Separately, we fit an ensemble model of distribution families to describe each CGF Z score distribution from individual-level data. Ten distributions were fit simultaneously (normal, log-normal, log-logistic, exponential, gamma, mirrored gamma, inverse gamma, Gumbel, mirrored Gumbel, and Weibull). We derived weights for each distribution based on minimised error in predicting CGF prevalence of less than -1 Z score, less than -2 Z scores, and less than -3 Z scores. Finally, we synthesised the results from these steps to estimate continuous HAZ, WAZ, and WHZ curves to estimate a weighted probability density function based on the ensemble distributions and ST-GPR mean and SD values.

Estimation of the relative risk of cause-specific incidence and mortality at varying levels of CGF indicators

We created new estimates of the relative risk of cause-specific incidence and mortality for continuous distributions of CGF exposure for GBD 2021 and used these estimates for GBD 2023. Estimates of CGF burden in previous GBD publications used categorical exposures (-3 , -2 , and -1 Z scores) matched with relative risks of cause-specific disease burden at each of those levels of exposure.^{22,23} A longer description of the methods is available elsewhere,²² and we have summarised the methods below.

For cause-specific mortality, we included 22 870 children and more than 137 000 anthropometric measurements from eight studies, representing 1029 cause-specific deaths (283 due to diarrhoea, 737 due to acute lower respiratory infections, and nine due to malaria). We also included hazard ratios for cause-specific mortality from a pooled analysis by Olofin and colleagues³ in our meta-regression. For cause-specific incidence, we included 60 436 children and more than 286 000 anthropometric measurements from 19 studies, representing more than 100 000 unique study-defined infectious disease episodes (95 950 diarrhoea episodes, 12 947 acute lower respiratory infections, 4896 malaria episodes, and 376 measles episodes). These data were from longitudinal studies that reported multiple measurements of height and bodyweight over time. We collapsed these observations into bins of Z scores for each CGF indicator and by age and study while summing the number of child-days of infectious disease episodes based on a standardised one-week recall, and cause-specific deaths within each bin.

We used a Bayesian meta-regression tool and risk factor assessment framework built for GBD and used across risk factors to quantify a continuous risk curve for each CGF exposure category and outcome.²⁴ Each risk exposure indicator (HAZ, WAZ, and WHZ) across the Z score range -6 to 1 was paired with each outcome (incidence and mortality for each infectious disease) to create a risk-outcome pair. In other words, we built separate models for the incidence and mortality rates for each infectious disease outcome, and for each metric of growth failure (24 models). We used logit-transformed mortality ratio and log-transformed incidence as the continuous independent variables in our models. Age was included as a fixed effect in the model. Studies were included as random intercepts in the model, and a flexible regularised spline fitting algorithm determined the shape of the curves. Age was chosen as a fixed effect since we wished to estimate the specific impact of age on the outcomes while our random effects were chosen as such due to our interest in estimating both the differences by study as well as the general variation one would expect from study to study. The meta-regression model incorporates both within-study and between-study uncertainty measured by standard errors and simulates Bayesian uncertainty intervals including and excluding between-study variation. More details on the meta-regression tool can be found in the publication associated with the method²⁴ as well as the publication from which the relative risk estimates are taken.²³

Because of the high degree of correlation between stunting, underweight, and wasting, we adjusted our relative risk values by simulating a joint distribution of the three indicators using extracted data from Demographic and Health Surveys. Based on an analysis done by McDonald and colleagues,⁵ we fit an interaction term between the three indicators and calculated adjusted relative risks by minimising the error between the

crude relative risks from our meta-regression and the expected relative risk derived from the joint estimate with the interaction term. More details of the specific approach taken are provided in appendix 1 of the report by the GBD 2023 Disease and Injury and Risk Factor Collaborators.²²

Calculation of a cause-age-sex-location-specific attributable fraction and multiplication by cause burden

Given that there were two continuous distributions, we produced estimates of the attributable fraction of disease incidence and mortality. The attributable fraction represents the proportion of disease burden that was attributable to a given risk factor (ie, the proportion of disease burden that would not exist in the absence of the risk factor). As CGF does not directly cause infections or deaths, the interpretation of the percentage attributable fraction is the proportion of disease burden that would not exist in the absence of the risk factor due to the linked reduction in associated direct biological drivers that cause increased burden such as immune system dysregulation.

The attributable fraction for continuous population-level exposure estimates and relative risks is:

$$PAF_{i,m,c} = \frac{\int_0^{\infty} RR_{i,m,c}(x) P_i(x) dx - 1}{\int_0^{\infty} RR_{i,m,c}(x) P_i(x) dx}$$

where i is the CGF indicator (HAZ, WAZ, or WHZ), m is the disease outcome (incidence or death), and c is the cause (diarrhoea, lower respiratory infections, malaria, or measles). An attributable fraction for CGF overall was estimated with a multiplicative aggregation of the individual indicator attributable fractions:

$$PAF_{m,c} = 1 - \prod_{i=1}^n (1 - PAF_{i,m,c})$$

Notably, due to a scarcity of data, GBD does not estimate attributable burden for stunting in the early neonatal (0–6 days) and late neonatal (7–27 days) age groups. Attribution is estimated for the following age groups: 1 to less than 6 months, 6 to less than 12 months, 12 to less than 24 months, 24 to less than 60 months.

Cause-specific incidence and mortality modelling

A full description of the modelling methods for the incidence and mortality of diarrhoea, lower respiratory infections, malaria, and measles has been published previously.^{25,26}

In short, mortality due to diarrhoea, lower respiratory infections, and measles was modelled with a Bayesian ensemble hierarchical tool called the Cause of Death Ensemble model (CODEm).²⁷ Measles used CODEm for settings with vital registration systems and used a separate model based on incidence and case-fatality ratio

for other settings.²⁷ Malaria mortality was modelled with spatially defined incidence and case-fatality estimates.^{27,28} Input data included vital registration, verbal autopsy, administrative records, and surveys. Covariates informed the models, and the final set of ensemble models was selected on the basis of out-of-sample predictive validity. Non-fatal incidence of diarrhoea and lower respiratory infections was modelled in a meta-regression tool with data from population-representative surveys, clinical data, administrative data, and scientific literature.²⁷ This meta-regression tool includes a compartmental

component that enforces consistency between disease incidence and mortality, making it an internally consistent estimate with mortality. Malaria incidence was based on estimates from the Malaria Atlas Project and on administrative, routine surveillance, and other geolocated and community-representative observations of infection prevalence for *Plasmodium falciparum*.²⁸ Measles incidence was estimated in a mixed-effects regression model by use of Joint Reporting Form case notifications and 5-year rolling lagged routine measles vaccination rates.²⁷

	Child growth failure	Child underweight	Child wasting	Child stunting
Global				
All causes	79 400 000 (47 000 000 to 106 000 000)	52 200 000 (21 900 000 to 75 100 000)	39 200 000 (23 800 000 to 53 000 000)	33 000 000 (24 100 000 to 42 200 000)
Diarrhoeal diseases	21 800 000 (13 000 000 to 32 200 000)	10 600 000 (5 870 000 to 17 400 000)	15 100 000 (-1 090 000 to 29 200 000)	7 580 000 (4 290 000 to 11 700 000)
Lower respiratory infections	32 100 000 (22 400 000 to 41 200 000)	22 100 000 (16 900 000 to 38 000 000)	12 500 000 (7 900 000 to 18 100 000)	17 300 000 (12 200 000 to 23 400 000)
Malaria	9 600 000 (-6 360 000 to 30 400 000)	6 980 000 (-4 070 000 to 20 500 000)	..	4 460 000 (-2 350 000 to 19 700 000)
Measles	6 930 000 (2 700 000 to 13 300 000)	3 560 000 (1 330 000 to 6 880 000)	2 590 000 (782 000 to 5 530 000)	3 690 000 (1 320 000 to 7 100 000)
Central Europe, eastern Europe, and central Asia				
All causes	792 000 (577 000 to 1 030 000)	442 000 (54 500 to 876 000)	314 000 (246 000 to 393 000)	303 000 (198 000 to 406 000)
Diarrhoeal diseases	70 500 (32 900 to 105 000)	22 000 (12 500 to 34 800)	51 000 (-2120 to 97 000)	17 900 (11 800 to 25 600)
Lower respiratory infections	706 000 (526 000 to 920 000)	405 000 (15 000 to 841 000)	248 000 (158 000 to 342 000)	285 000 (184 000 to 385 000)
Malaria	0 (0 to 0)	0 (0 to 0)	..	0 (0 to 0)
Measles	1040 (501 to 1450)	351 (163 to 510)	374 (140 to 751)	510 (233 to 737)
High income				
All causes	57 900 (37 100 to 92 300)	30 100 (9950 to 64 800)	24 800 (18 200 to 32 900)	17 700 (13 900 to 23 500)
Diarrhoeal diseases	11 500 (2420 to 23 200)	2750 (1840 to 4370)	7450 (-436 to 19 100)	2450 (1130 to 3840)
Lower respiratory infections	38 800 (26 600 to 64 000)	19 800 (514 to 54 600)	9830 (5620 to 14 400)	15 200 (10 900 to 21 100)
Malaria	0.0753 (-0.0302 to 0.312)	0.0753 (-0.0302 to 0.312)	..	<0.001 (->0.001 to <0.001)
Measles	4.48 (1.66 to 6.86)	1.25 (0.478 to 2.01)	1.77 (0.561 to 3.58)	1.73 (0.654 to 2.83)
Latin America and Caribbean				
All causes	1 470 000 (1 110 000 to 1 850 000)	920 000 (427 000 to 1 430 000)	624 000 (450 000 to 788 000)	566 000 (417 000 to 728 000)
Diarrhoeal diseases	322 000 (202 000 to 460 000)	118 000 (67 800 to 188 000)	183 000 (-6980 to 415 000)	115 000 (76 600 to 159 000)
Lower respiratory infections	878 000 (641 000 to 1 180 000)	528 000 (22 700 to 1 070 000)	168 000 (97 600 to 242 000)	451 000 (318 000 to 578 000)
Malaria	1810 (-892 to 6850)	1320 (-665 to 4290)	..	684 (-239 to 3520)
Measles	3.5 (1.53 to 5.11)	1.09 (0.494 to 1.64)	0.849 (0.322 to 1.76)	1.95 (0.797 to 2.86)
North Africa and Middle East				
All causes	2 800 000 (2 020 000 to 3 590 000)	1 790 000 (884 000 to 2 690 000)	1 490 000 (1 070 000 to 1 920 000)	1 120 000 (816 000 to 1 470 000)
Diarrhoeal diseases	582 000 (284 000 to 1 010 000)	254 000 (128 000 to 458 000)	403 000 (-20 700 to 873 000)	203 000 (107 000 to 345 000)
Lower respiratory infections	1 480 000 (1 010 000 to 2 050 000)	947 000 (61 500 to 1 830 000)	565 000 (328 000 to 848 000)	749 000 (486 000 to 1 060 000)
Malaria	43 800 (-33 200 to 126 000)	34 500 (-23 800 to 98 100)	..	17 600 (-7240 to 72 500)
Measles	282 000 (99 700 to 585 000)	138 000 (50 100 to 288 000)	103 000 (33 100 to 217 000)	154 000 (50 100 to 321 000)
South Asia				
All causes	15 400 000 (12 500 000 to 18 100 000)	10 800 000 (5 050 000 to 14 600 000)	9 500 000 (7 040 000 to 11 500 000)	6 210 000 (4 460 000 to 8 120 000)
Diarrhoeal diseases	4 020 000 (2 660 000 to 5 630 000)	2 180 000 (1 400 000 to 3 290 000)	2 930 000 (-263 000 to 5 150 000)	1 390 000 (875 000 to 2 060 000)
Lower respiratory infections	8 790 000 (6 880 000 to 10 700 000)	6 310 000 (526 000 to 9 610 000)	4 430 000 (3 020 000 to 6 100 000)	4 430 000 (2 850 000 to 6 110 000)
Malaria	179 000 (-145 000 to 598 000)	140 000 (-97 700 to 450 000)	..	76 400 (-37 100 to 360 000)
Measles	602 000 (231 000 to 1 270 000)	336 000 (119 000 to 717 000)	285 000 (84 700 to 623 000)	305 000 (111 000 to 653 000)

(Table 1 continues on next page)

	Child growth failure	Child underweight	Child wasting	Child stunting
(Continued from previous page)				
Southeast Asia, east Asia, and Oceania				
All causes	3 730 000 (2 810 000 to 4 660 000)	2 380 000 (1 020 000 to 3 520 000)	1 840 000 (1 320 000 to 2 370 000)	1 520 000 (1 110 000 to 1 990 000)
Diarrhoeal diseases	924 000 (472 000 to 1 530 000)	417 000 (208 000 to 751 000)	611 000 (-33 600 to 1 340 000)	322 000 (168 000 to 552 000)
Lower respiratory infections	2 130 000 (1 500 000 to 2 800 000)	1 430 000 (96 800 to 2 580 000)	737 000 (439 000 to 1 070 000)	1 060 000 (703 000 to 1 480 000)
Malaria	12 700 (-8810 to 44 300)	9450 (-5840 to 30 700)	..	5450 (-2700 to 26 400)
Measles	246 000 (91 200 to 486 000)	118 000 (41 700 to 245 000)	82 000 (24 600 to 172 000)	133 000 (49 500 to 273 000)
Sub-Saharan Africa				
All causes	55 200 000 (26 700 000 to 77 100 000)	35 800 000 (13 400 000 to 53 100 000)	25 400 000 (14 400 000 to 36 800 000)	23 300 000 (14 600 000 to 34 000 000)
Diarrhoeal diseases	15 900 000 (9 100 000 to 24 400 000)	7 600 000 (4 100 000 to 12 600 000)	10 900 000 (-762 000 to 21 600 000)	5 530 000 (2 950 000 to 8 790 000)
Lower respiratory infections	18 100 000 (11 100 000 to 25 400 000)	12 400 000 (969 000 to 22 400 000)	6 360 000 (3 670 000 to 10 200 000)	10 300 000 (6 660 000 to 14 400 000)
Malaria	9 360 000 (-6 160 000 to 29 600 000)	6 790 000 (-3 940 000 to 20 000 000)	..	4 360 000 (-2 290 000 to 19 200 000)
Measles	5 800 000 (2 210 000 to 11 100 000)	2 960 000 (1 090 000 to 5 690 000)	2 120 000 (628 000 to 4 460 000)	3 100 000 (1 070 000 to 6 070 000)
Estimates combine the burden associated with mild, moderate, and severe forms of child growth failure: stunting was defined as height-for-age Z score (HAZ) less than -1; underweight was defined as weight-for-age Z score (WAZ) less than -1; and wasting was defined as weight-for-height Z score (WHZ) less than -1, according to WHO Child Growth Standards. The meta-regression model used to estimate relative risks for various levels of risk-factor exposure incorporates both within-study and between-study uncertainty measured by standard errors. The results presented herein incorporate between-study variation, meaning the uncertainty intervals here represent both uncertainty in the relative risks as well as predicted variation due to between-study variation, and as such some uncertainty intervals' lower bounds fall below zero. As described in the publication associated with the method, ²⁴ a risk factor, outcome, or cause combination would not be included in the table if the uncertainty without between-study variation was not statistically significantly different from zero. Data in parentheses are 95% uncertainty intervals. Count data are presented to three significant figures. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.				
Table 1: All-cause and cause-specific DALYs associated with child growth failure among children younger than 5 years at the global and GBD super-regional levels, 2023				

Cause-specific cases and deaths were multiplied by the population attributable fraction (PAF) values to produce our final estimates of the disease burden associated with each CGF indicator. All steps in the estimation process included 500 iterations of each age-sex-geography-year-specific value. 95% uncertainty intervals (UIs) were calculated as the 2.5th and 97.5th percentiles across all draws. This uncertainty was carried through the entire process to maintain the variation in the estimates. Count data are presented to three significant figures, while rates and percentages are presented to one decimal place.

The code and input data are available on the Global Health Data Exchange (GHDx) website. This study complies with the GATHER requirements for burden estimation and reporting.²⁹

Role of the funding source

The funder of this study had no role in study design, data collection, data analysis, data interpretation, or the writing of the report.

Results

Globally, in 2023, in children younger than 5 years, all CGF (mild, moderate, and severe) was associated with 79.4 million (95% UI 47.0–106) DALYs, ranking only behind low birthweight and short gestation (133 million [125–143] DALYs) as the leading risk factors associated with health burden among all children younger than 5 years (table 1; appendix 1 table S1). CGF was associated with 17.9% (10.6–23.8) of 444 million (434–457) total under-5 DALYs (table 2; appendix table S2). In 2023, CGF

was associated with 880 000 (517 000–1 170 000) deaths (figures 1, 2; table 3; appendix 1 table S3), representing 18.8% (11.1–25.0) of all 4.67 million (4.59–4.75) under-5 deaths (figure 1; table 4; appendix 1 table S4) and 1.54 million (0.804–2.41) years lived with disability (YLDs; appendix 1 table S5), representing 5.5% (3.2–7.9) of all 28.0 million (20.4–37.1) under-5 YLDs (figure 2; appendix 1 table S6). Due to limited space and similar trends between DALYs and mortality through time and across countries by indicator and cause, we primarily focus the results presented below on mortality. Full estimates produced in association with these analyses across all health metrics are available via the GBD Results Tool.

The number of deaths associated with CGF was 2.75 million (95% UI 1.95–3.27) in 2000, decreasing by an average of 5.0% (4.2–5.9) per year to 880 000 (517 000–1 170 000) in 2023 (figure 1; GBD Results Tool). Some of these decreases were due to secular trends in disease burden not related to CGF. The attributable fraction of CGF decreased at a slower rate, by 2.1% (1.4–3.0) per year since 2000, from 30.5% (21.5–36.3) of deaths in children younger than 5 years in 2000 to 18.8% (11.1–25.0) in 2023 (figure 1; table 4). However, the attributable fraction of CGF among all causes of death for which it contributes as a risk factor decreased less rapidly between 2000, from 73.7% (53.9–86.3) in 2000 to 61.2% (33.6–79.2) in 2023, representing a 0.9% (0.3–2.0) average yearly reduction (GBD Results Tool).²²

Among the different indicators of CGF, underweight was associated with the greatest disease burden:

For the code and input data see <https://ghdx.healthdata.org/gbd-2023/code>

For the GBD Results Tool see <https://vizhub.healthdata.org/gbd-results/>

12·3% (95% UI 5·1–17·6) of all deaths in children younger than 5 years in 2023, followed by child wasting (9·2% [5·5–12·5]) and child stunting (8·0% [6·0–10·3]; figure 1, table 4; appendix 1 table S4). The fastest decline in attributable burden from 2000 was in child stunting (2·6% [1·8–3·6] decline per year), followed by child wasting (2·4% [1·5–3·2] decline per year) and underweight (2·3% [1·5–3·2] decline per year; figure 1).

In 2023, at the GBD super-regional level, the highest percentage of total under-5 deaths associated with CGF occurred in sub-Saharan Africa (23·4% [95% UI 11·3 to 32·8]) and in south Asia (14·6% [11·8 to 16·8]; table 4; appendix 1 table S4). In 2023, CGF was associated with 618 000 (299 000 to 862 000) deaths in sub-Saharan Africa and 165 000 (134 000 to 194 000) deaths in south Asia (table 3, appendix 1 table S3). At the national level, in 2023, the largest numbers

	Child growth failure	Child underweight	Child wasting	Child stunting
Global				
All causes	17·9% (10·6 to 23·8)	11·7% (5·0 to 16·8)	8·8% (5·4 to 11·9)	7·4% (5·6 to 9·5)
Diarrhoeal diseases	75·2% (53·1 to 88·1)	36·6% (24·9 to 47·3)	51·9% (-3·1 to 84·2)	26·2% (18·6 to 31·9)
Lower respiratory infections	59·1% (48·5 to 66·5)	40·5% (2·7 to 63·0)	23·0% (16·0 to 29·3)	31·8% (24·3 to 37·9)
Malaria	26·3% (-17·8 to 69·4)	19·1% (-11·6 to 48·9)	..	12·1% (-6·2 to 44·7)
Measles	66·4% (39·9 to 75·8)	34·0% (18·3 to 42·4)	24·9% (10·8 to 40·9)	35·3% (17·3 to 43·5)
Central Europe, eastern Europe, and central Asia				
All causes	11·2% (8·2 to 14·7)	6·2% (0·76 to 12·6)	4·4% (3·5 to 5·5)	4·3% (2·8 to 5·7)
Diarrhoeal diseases	56·4% (27·8 to 78·1)	17·7% (10·5 to 24·7)	40·7% (-1·8 to 74·2)	14·4% (10·0 to 18·3)
Lower respiratory infections	43·5% (32·8 to 57·2)	24·9% (0·93 to 52·3)	15·3% (10·0 to 20·4)	17·5% (11·4 to 23·5)
Malaria	0% (0 to 0)	0% (0 to 0)	..	0% (0 to 0)
Measles	38·9% (18·9 to 48·4)	13·1% (6·2 to 17·5)	14·0% (5·3 to 25·6)	19·0% (8·7 to 25·4)
High income				
All causes	0·98% (0·62 to 1·6)	0·51% (0·17 to 1·1)	0·42% (0·30 to 0·57)	0·30% (0·23 to 0·41)
Diarrhoeal diseases	18·4% (3·9 to 35·5)	4·4% (2·8 to 6·9)	11·9% (-0·69 to 29·1)	3·9% (1·7 to 6·1)
Lower respiratory infections	34·6% (24·4 to 54·5)	17·6% (0·44 to 47·2)	8·8% (5·1 to 12·2)	13·6% (9·9 to 17·7)
Malaria	8·3% (-3·4 to 29·8)	8·3% (-3·4 to 29·8)	..	<0·1% (>-0·1 to <0·1)
Measles	13·6% (6·0 to 17·7)	3·8% (1·8 to 5·2)	5·4% (1·9 to 10·0)	5·2% (2·4 to 7·2)
Latin America and Caribbean				
All causes	9·2% (6·9 to 11·6)	5·8% (2·7 to 9·1)	3·9% (2·8 to 5·0)	3·5% (2·6 to 4·5)
Diarrhoeal diseases	55·0% (36·4 to 73·7)	20·2% (12·1 to 28·7)	31·3% (-1·1 to 66·5)	19·6% (13·7 to 24·7)
Lower respiratory infections	52·2% (38·7 to 68·7)	31·3% (1·3 to 63·2)	10·0% (6·0 to 13·6)	26·8% (18·7 to 33·5)
Malaria	14·5% (-7·2 to 45·5)	10·8% (-5·3 to 31·2)	..	5·1% (-2·2 to 23·4)
Measles	25·7% (12·3 to 32·4)	8·0% (3·8 to 10·6)	6·2% (2·4 to 11·7)	14·3% (6·5 to 19·7)
North Africa and Middle East				
All causes	10·1% (7·3 to 12·8)	6·5% (3·2 to 9·6)	5·4% (3·9 to 6·9)	4·1% (3·0 to 5·3)
Diarrhoeal diseases	64·7% (39·9 to 80·9)	28·2% (19·2 to 37·9)	44·7% (-2·1 to 77·1)	22·5% (16·2 to 27·9)
Lower respiratory infections	54·0% (43·8 to 63·1)	34·5% (2·0 to 58·7)	20·7% (13·9 to 27·0)	27·5% (18·9 to 34·8)
Malaria	31·3% (-23·3 to 78·1)	24·7% (-17·2 to 60·2)	..	12·3% (-6·3 to 49·0)
Measles	61·1% (34·4 to 70·9)	29·8% (15·1 to 38·1)	22·4% (9·3 to 38·7)	33·3% (15·5 to 41·2)
South Asia				
All causes	14·1% (11·4 to 16·3)	9·9% (4·6 to 12·8)	8·7% (6·4 to 10·2)	5·7% (4·1 to 7·3)
Diarrhoeal diseases	77·1% (55·2 to 86·8)	41·8% (30·2 to 51·9)	56·2% (-3·9 to 83·6)	26·7% (19·0 to 32·4)
Lower respiratory infections	53·1% (45·1 to 58·8)	38·2% (2·9 to 55·9)	26·8% (19·4 to 33·3)	26·8% (17·3 to 33·8)
Malaria	32·1% (-24·4 to 78·9)	25·2% (-17·7 to 60·4)	..	13·4% (-7·2 to 50·9)
Measles	75·2% (48·7 to 84·8)	42·0% (23·7 to 51·0)	35·8% (16·4 to 55·2)	38·0% (19·0 to 46·4)
Southeast Asia, east Asia, and Oceania				
All causes	11·2% (8·4 to 13·7)	7·1% (3·0 to 10·4)	5·5% (4·0 to 7·1)	4·6% (3·4 to 5·9)
Diarrhoeal diseases	68·5% (44·7 to 83·9)	31·0% (21·3 to 40·3)	45·1% (-2·3 to 78·0)	23·8% (17·2 to 29·2)
Lower respiratory infections	56·5% (43·5 to 66·0)	37·8% (2·2 to 62·6)	19·5% (13·0 to 25·3)	28·0% (19·3 to 34·9)
Malaria	26·1% (-18·5 to 69·2)	19·6% (-12·5 to 51·3)	..	10·6% (-5·5 to 42·1)
Measles	60·9% (35·4 to 70·2)	29·3% (15·6 to 36·7)	20·5% (8·5 to 35·1)	32·9% (16·3 to 40·7)

(Table 2 continues on next page)

	Child growth failure	Child underweight	Child wasting	Child stunting
(Continued from previous page)				
Sub-Saharan Africa				
All causes	22.5% (10.9 to 31.7)	14.6% (5.5 to 21.7)	10.4% (5.9 to 15.0)	9.5% (6.0 to 13.9)
Diarrhoeal diseases	76.5% (54.2 to 89.5)	36.6% (24.5 to 47.9)	52.4% (-3.0 to 86.0)	26.6% (18.9 to 32.7)
Lower respiratory infections	64.8% (52.2 to 73.6)	44.4% (2.9 to 69.3)	22.8% (15.1 to 30.0)	36.9% (30.2 to 42.9)
Malaria	26.2% (-17.7 to 69.3)	19.0% (-11.5 to 48.7)	..	12.1% (-6.2 to 44.6)
Measles	66.1% (39.4 to 75.7)	33.8% (18.1 to 42.1)	24.3% (10.4 to 40.2)	35.3% (17.3 to 43.6)
<p>Estimates combine the burden associated with mild, moderate, and severe forms of child growth failure: stunting was defined as height-for-age Z score (HAZ) less than -1; underweight was defined as weight-for-age Z score (WAZ) less than -1; and wasting was defined as weight-for-height Z score (WHZ) less than -1, according to WHO Child Growth Standards. The results presented herein incorporate between-study variation, meaning the uncertainty intervals here represent both uncertainty in the relative risks as well as predicted variation due to between-study variation, and as such some uncertainty intervals' lower bounds fall below zero. As described in the publication associated with the method,²⁴ a risk factor, outcome, or cause combination would not be included in the table if the uncertainty without between-study variation was not statistically significantly different from zero. Data in parentheses are 95% uncertainty intervals. Population attributable fractions are presented to one decimal place. DALY=disability-adjusted life-year. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.</p>				
<p>Table 2: All-cause and cause-specific population attributable fraction of DALYs among children younger than 5 years for child growth failure at the global and GBD super-regional levels, 2023</p>				

of deaths among children younger than 5 years associated with CGF occurred in Nigeria (188 000 [70 200 to 270 000]), India (112 000 [88 300 to 131 000]), and DR Congo (50 800 [17 000 to 87 800]; appendix 1 table S3). The highest proportions of under-5 deaths associated with CGF in 2023 were in Chad (40.2% [27.9 to 48.8] of all under-5 deaths), Niger (34.9% [17.3 to 45.5]), and South Sudan (32.8% [18.9 to 43.3]; appendix 1 table S4). Globally, in 2023, CGF was associated with 362 000 (253 000 to 465 000) deaths due to lower respiratory infections in children younger than 5 years (59.4% [48.9 to 66.8] of 610 000 [486 000 to 749 000] deaths due to lower respiratory infection in children <5 years), followed by diarrhoeal diseases (243 000 [146 000 to 360 000]; 76.7% [54.9 to 89.1] of 317 000 [213 000 to 462 000] under-5 diarrhoea deaths), malaria (107 000 [-71 300 to 341 000]; 26.4% [-17.8 to 69.5] of 409 000 [158 000 to 735 000] under-5 malaria deaths), and measles (78 600 [30 700 to 150 000] deaths; 66.6% [40.0 to 76.0] of 118 000 [47 700 to 211 000] under-5 measles deaths; tables 3, 4; appendix 1 figures S1–S5; tables S3, S4). CGF was also associated with the burden of cause-specific infectious disease incidence, including 27.7% (-24.4 to 70.7) of 605 000 (387 000 to 884 000) under-5 malaria YLDs, 24.7% (-18.1 to 56.9) of 767 000 (527 000 to 1 070 000) under-5 diarrhoeal YLDs, 13.4% (-32.7 to 50.4) of 60 000 (40 100 to 85 900) under-5 lower respiratory infection YLDs, and 11.7% (-4.9 to 30.3) of 38 400 (15 000 to 80 000) under-5 measles YLDs (appendix 1 figures S1–S4, table S6). The CGF-attributable burden by cause varied substantially between countries and regions (figure 3; appendix 1 figure S5). The attributable fraction by cause was lowest among countries in the high-income super-region, but the range of deaths due to diarrhoea and lower respiratory infection associated with CGF still varied from 6.0% to 51.2% between countries in this super-region (appendix 1 table S2).

Discussion

In 2023, CGF was a leading factor associated with mortality in children younger than 5 years globally, falling only behind short gestation and low birthweight, and was associated with approximately 791 000 deaths due to diarrhoea, lower respiratory infections, malaria, and measles. The burden associated with CGF decreased between 2000 and 2023 and has been an important correlate for tracking reductions in childhood mortality.^{30,31} Multifaceted strategies are needed to address the intersectional contribution of environmental, socioeconomic, biological, and behavioural factors that affect childhood growth.

No single intervention is likely to improve childhood growth for all children.¹ Recent reviews of effective interventions for maternal and childhood malnutrition describe interventions (appendix 1 table S7) according to direct and indirect relationships between the health sector or other macro-level sectors and undernutrition and provide a useful framework to consider different interventions at different points in the life cycle.^{32,33} Although societal and indirect interventions, such as family planning and reproductive health services, or poverty alleviation and women's empowerment strategies, might have important effects on CGF, several reviews have focused on the more proximal interventions that affect childhood environment and biology.^{32,34}

Across the CGF indicators, childhood underweight was associated with the largest proportion of the under-5 disease burden. Due to the calculations for underweight, wasting, and stunting, WAZ scores are depressed by both wasting and stunting, making WAZ lowest when both substantial wasting and stunting are present. Moreover, there is evidence that wasting and stunting reinforce each other, meaning that stunting increases the risk of future wasting and vice versa—a relationship that gets stronger as a child grows older.¹² As such, locations with a high estimated burden




Figure 1: Global burden associated with child growth failure for children younger than 5 years as a fraction of all deaths (A), number of attributable deaths (B), and attributable mortality rate per 100 000 (C) Separate panels show the attributable burden for underweight, wasting, and stunting separately, as well as child growth failure as a whole. Shaded regions represent 95% uncertainty intervals. Estimates combine burden associated with mild, moderate, and severe forms of child growth failure: stunting was defined as height-for-age Z score (HAZ) less than -1; underweight was defined as weight-for-age Z score (WAZ) less than -1; and wasting was defined as weight-for-height Z score (WHZ) less than -1, according to WHO Child Growth Standards.

associated with all three CGF indicators are likely to require disproportionately more intense interventions than those where stunting and wasting are reinforcing each other.

Recovery from stunting is rare: most children who experience stunting do not recover towards the global mean for length or height, and those who do gain a small amount on their peers.³⁵ Although there is some evidence that stunting can be addressed after the first 1000 days of life,³⁶ early detection and intervention are critical for successful reversal of stunting.⁴ Our work found that a larger proportion of the CGF burden was associated with stunting than in previous estimates; given the difficulty in reversing stunting, our estimates should be used to identify locations in need of increased early detection and intervention. In a pooled analysis by Mertens and colleagues,¹² comprising 19 studies and about 60 000 infants, the cumulative incidence, including at birth, of stunting was 25% in the first 3 months of life.¹² This study also found that the incidence of wasting peaked in the first 3 months of life.¹² Recovery from wasting is more common than from stunting. In the pooled analysis,¹² about 65% of infants recovered from wasting within 60 days in the first 3 months of life, and about 50% of infants who experienced wasting at older ages recovered in 60 days. This suggests that changes in lifestyle or interventions have a larger impact in younger compared to older children and for wasting compared to stunting.

Early detection must begin before birth. On average, children in low-income and middle-income settings tend to grow more slowly than the WHO global reference standard, moving further away from the global mean as they age up to 5 years. In part, this appears to be because they are born small, and children born with low birthweight or of short gestation are more likely to experience growth failure.¹⁴ Other reviews support the importance of intervening in the pre-conception, pregnancy, or neonatal periods to prevent the burden of CGF.^{14,15,32}

There are several potential mechanisms that explain how suboptimal growth is associated with an increased risk of infectious disease burden, including an altered microbiome, chronic inflammation, immune system dysregulation, changes in endothelial barrier function, and others.³⁷ A review of the literature concluded that although improvements in nutrition can reverse susceptibility to infection, this effect is not well characterised.¹ Reversing acute malnutrition might help make children more resilient to infection.

Our study includes meaningful updates to previously reported estimates of the CGF burden in GBD 2019.² We used continuous estimates of height and bodyweight by age, sex, geography, and year instead of modelled estimates of the prevalence of CGF indicators. This approach reflects the continuous nature of deviations in growth from an expected mean. We also used continuous

estimates of the relative risks of incidence and mortality due to infectious causes for each CGF indicator by age group. Together, these improvements have produced estimates of a higher specificity and with a better reflection of the true, underlying relationship between population-level exposures and risk factors.

Since GBD 2021, updated relative risk estimates have been used to quantify the burden of disease that is associated with CGF.²² Previous iterations of GBD (eg, GBD 2017 and GBD 2019)² used the cause-specific mortality hazard ratios from the study by Olofin and colleagues³ for both incidence and mortality. In addition to estimating the risk of cause-specific disease incidence, there are two main ways in which these estimates differ from those values. First, the relative risk of lower respiratory infection due to low HAZ and low WAZ are meaningfully higher than in the study by Olofin and colleagues³ and in comparison with a separate systematic review and network meta-analysis.^{22,38} Second, we identified statistically significant relationships between low HAZ and low WAZ and malaria mortality, which were not observed in the smaller sample size and discrete categorical analysis by Olofin and colleagues.³ These CGF indicators were only included as risk factors associated with malaria mortality in GBD 2021 based on these findings and might therefore represent a gap in previous estimates of the malaria burden.

Although our updated CGF burden estimates highlight changes in the proportions of the under-5 health burden associated with wasting, stunting, and underweight compared to previous iterations of GBD, the total CGF burden remains comparable to that of GBD 2021, GBD 2019, and GBD 2017. However, even with increased relative risks and estimates of the stunting burden and after adjusting for reductions in the CGF burden in the past decade, our stunting estimates remain considerably lower than those of the Maternal and Child Nutrition Study Group (MCNS).⁴ The MCNS group used CGF indicator prevalence estimated by the UN³⁹ and the Nutrition Impact Model Study⁴⁰ as well as cause-specific relative risks from a pooled analysis of ten longitudinal studies³ to estimate CGF-attributable deaths in children younger than 5 years in 2011 (appendix 1 table S8). Our estimates are based on aggregating the burden associated with mild, moderate, and severe forms of each CGF indicator (eg, Z scores <-1) while the MCNS group focused only on moderate and severe forms (eg, Z scores <-2). In spite of this, our underweight and wasting burden estimates are somewhat comparable with those of the MCNS group. A comprehensive comparison between estimates would require comparing the burden using identical definitions of stunting, wasting, and underweight, which is difficult considering that it depends on exposure prevalence, risk of cause-specific death, and the total number of under-5 and cause-specific deaths. Importantly, no other group, to date, has used the updated relative risk estimates for




Figure 2: Percentage of all-cause deaths and years lived with disability associated with all child growth failure (mild, moderate, and severe) in children younger than 5 years, globally, in 2023

Population attributable fraction of all-cause deaths and years lived with disability for all children younger than 5 years, as well as age-specific values (1 to <6 months, 6 to <12 months, 12 to <24 months, and 24 to <60 months) are plotted. Dots represent mean population attributable fractions and whiskers represent 95% uncertainty intervals.

CGF indicators used in the current study, which probably explains some of the differences in estimates.

There are several limitations to this analysis. First, our estimates depend on numerous sources of data for childhood growth, infectious disease incidence, and infectious disease mortality, each with various potential gaps and biases, which we attempted to resolve using expert opinion and sophisticated statistical models. One common data gap for each of the inputs into this study is that the burden of CGF is highest in countries in the lowest SDI quintiles that do not have robust vital registration and disease surveillance systems. We attempted to account for uncertainty throughout our modelling process by including measured error when possible and producing estimates incorporating uncertainty in each step. We strived to report our findings both as the mean from the posterior distributions as well as the 2.5th and 97.5th percentiles of those distributions. Capturing and reporting uncertainty is crucial in modelling burden of disease, including this report. Second, many unanswered questions remain about the impact of the COVID-19 pandemic and the associated disruptions in maternal and child health services on child nutrition and mortality.⁴¹ We have included modelled impacts on cause-specific child mortality,⁴² but strong evidence of changes in vaccine coverage or birth size and gestational age, two confounding risks for infectious disease mortality in

	Child growth failure	Child underweight	Child wasting	Child stunting
Global				
All causes	880 000 (517 000 to 1 170 000)	573 000 (236 000 to 824 000)	428 000 (256 000 to 583 000)	373 000 (272 000 to 477 000)
Diarrhoeal diseases	243 000 (146 000 to 360 000)	118 000 (63 600 to 195 000)	169 000 (-11 900 to 327 000)	84 800 (49 900 to 132 000)
Lower respiratory infections	362 000 (253 000 to 465 000)	249 000 (19 200 to 428 000)	141 000 (88 700 to 204 000)	196 000 (138 000 to 264 000)
Malaria	107 000 (-71 300 to 341 000)	77 400 (-44 700 to 229 000)	..	50 800 (-26 800 to 224 000)
Measles	78 600 (30 700 to 150 000)	40 400 (15 200 to 78 100)	29 300 (8850 to 62 600)	42 000 (15 000 to 80 600)
Central Europe, eastern Europe, and central Asia				
All causes	8790 (6390 to 11 400)	4860 (512 to 9720)	3420 (2670 to 4320)	3410 (2230 to 4560)
Diarrhoeal diseases	774 (380 to 1140)	239 (123 to 386)	568 (-21.4 to 1080)	195 (117 to 284)
Lower respiratory infections	7920 (5900 to 10 300)	4540 (169 to 9430)	2770 (1770 to 3830)	3200 (2090 to 4320)
Malaria	0 (0 to 0)	0 (0 to 0)	..	0 (0 to 0)
Measles	11.7 (5.63 to 16.4)	3.96 (1.84 to 5.75)	4.18 (1.55 to 8.42)	5.76 (2.63 to 8.34)
High income				
All causes	623 (422 to 964)	322 (111 to 697)	269 (201 to 352)	193 (138 to 264)
Diarrhoeal diseases	107 (39.1 to 210)	19.6 (9.8 to 30.3)	78.2 (-1.7 to 202)	20.8 (12.5 to 29.5)
Lower respiratory infections	436 (299 to 718)	222 (5.93 to 612)	110 (63.1 to 162)	172 (124 to 238)
Malaria	<0.001 (>-0.001 to <0.001)	<0.001 (>-0.001 to <0.001)	..	<0.001 (>-0.001 to <0.001)
Measles	0.0505 (0.0185 to 0.0775)	0.0142 (0.00544 to 0.0228)	0.0198 (0.00615 to 0.0404)	0.0197 (0.00742 to 0.0321)
Latin America and Caribbean				
All causes	16 500 (12 500 to 20 800)	10 300 (4780 to 16 000)	6980 (5030 to 8820)	6370 (4660 to 8200)
Diarrhoeal diseases	3580 (2290 to 5100)	1310 (730 to 2110)	2060 (-75.7 to 4660)	1280 (824 to 1790)
Lower respiratory infections	9870 (7210 to 13 300)	5930 (256 to 12 000)	1890 (1090 to 2710)	5080 (3600 to 6500)
Malaria	19.2 (-9.6 to 74.3)	13.6 (-6.94 to 46.2)	..	7.78 (-2.72 to 40.0)
Measles	0.0394 (0.0172 to 0.0574)	0.0123 (0.0056 to 0.0186)	0.00933 (0.00331 to 0.0195)	0.022 (0.00896 to 0.0325)
North Africa and Middle East				
All causes	30 600 (21 900 to 39 400)	19 300 (9290 to 29 400)	15 900 (11 400 to 20 700)	12 600 (9160 to 16 600)
Diarrhoeal diseases	6410 (3260 to 11 100)	2780 (1340 to 5090)	4500 (-216 to 9740)	2240 (1150 to 3860)
Lower respiratory infections	16 600 (11 400 to 23 000)	10 600 (698 to 20 600)	6330 (3670 to 9520)	8440 (5490 to 11 900)
Malaria	472 (-357 to 1400)	366 (-253 to 1070)	..	200 (-82.7 to 827)
Measles	3200 (1130 to 6630)	1570 (568 to 3260)	1160 (373 to 2460)	1750 (568 to 3640)
South Asia				
All causes	165 000 (134 000 to 194 000)	113 000 (49 000 to 155 000)	98 900 (71 200 to 120 000)	69 600 (49 700 to 91 300)
Diarrhoeal diseases	44 200 (30 000 to 61 900)	23 800 (14 900 to 36 600)	32 600 (-2800 to 57 400)	15 400 (9690 to 23 100)
Lower respiratory infections	98 600 (77 200 to 120 000)	70 800 (5950 to 108 000)	49 600 (33 800 to 68 300)	49 900 (32 100 to 68 600)
Malaria	1990 (-1620 to 6700)	1540 (-1090 to 5020)	..	864 (-421 to 4070)
Measles	6810 (2620 to 14 300)	3810 (1350 to 8140)	3220 (956 to 7040)	3450 (1250 to 7410)
Southeast Asia, east Asia, and Oceania				
All causes	40 400 (30 300 to 50 500)	25 300 (10 200 to 37 400)	19 400 (13 800 to 25 400)	17 000 (12 400 to 22 400)
Diarrhoeal diseases	10 100 (5410 to 16 800)	4480 (2060 to 8360)	6800 (-339 to 14 900)	3520 (1690 to 6200)
Lower respiratory infections	24 000 (16 900 to 31 500)	16 000 (1100 to 28 900)	8260 (4920 to 12 100)	11 900 (7940 to 16 600)
Malaria	128 (-82.9 to 477)	90.6 (-51.0 to 317)	..	61.9 (-30.7 to 299)
Measles	2790 (1040 to 5520)	1350 (474 to 2790)	926 (277 to 1950)	1510 (562 to 3100)
Sub-Saharan Africa				
All causes	618 000 (299 000 to 862 000)	399 000 (149 000 to 594 000)	283 000 (160 000 to 412 000)	264 000 (165 000 to 386 000)
Diarrhoeal diseases	178 000 (104 000 to 274 000)	85 100 (45 200 to 143 000)	123 000 (-8480 to 243 000)	62 200 (33 400 to 99 100)
Lower respiratory infections	205 000 (126 000 to 288 000)	141 000 (11 000 to 253 000)	71 800 (41 300 to 115 000)	117 000 (75 700 to 163 000)
Malaria	105 000 (-69 000 to 333 000)	75 400 (-43 300 to 223 000)	..	49 700 (-26 200 to 219 000)
Measles	65 800 (25 100 to 127 000)	33 700 (12 400 to 64 700)	24 000 (7110 to 50 600)	35 200 (12 200 to 69 000)

Estimates combine burden associated with mild, moderate, and severe forms of child growth failure: stunting was defined as height-for-age Z score (HAZ) less than -1; underweight was defined as weight-for-age Z score (WAZ) less than -1; and wasting was defined as weight-for-height Z score (WHZ) less than -1, according to WHO Child Growth Standards. The results presented herein incorporate between-study variation, meaning the uncertainty intervals here represent both uncertainty in the relative risks as well as predicted variation due to between-study variation, and as such some uncertainty intervals' lower bounds fall below zero. As described in the publication associated with the method,²⁴ a risk factor, outcome, or cause combination would not be included in the table if the uncertainty without between-study variation was not statistically significantly different from zero. Data in parentheses are 95% uncertainty intervals. Count data are presented to three significant figures. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.

Table 3: All-cause and cause-specific deaths associated with child growth failure among children younger than 5 years at the global and GBD super-regional levels, 2023

	Child growth failure	Child underweight	Child wasting	Child stunting
Global				
All causes	18.8% (11.1 to 25.0)	12.3% (5.1 to 17.6)	9.2% (5.5 to 12.5)	8.0% (6.0 to 10.3)
Diarrhoeal diseases	76.7% (54.9 to 89.1)	37.2% (24.7 to 48.4)	53.2% (-3.1 to 86.1)	26.8% (19.2 to 33.0)
Lower respiratory infections	59.4% (48.9 to 66.8)	40.8% (2.7 to 63.3)	23.1% (16.0 to 29.4)	32.1% (24.7 to 38.1)
Malaria	26.4% (-17.8 to 69.5)	19.0% (-11.3 to 48.2)	..	12.4% (-6.4 to 45.9)
Measles	66.6% (40.0 to 76.0)	34.2% (18.4 to 42.5)	24.9% (10.8 to 41.0)	35.5% (17.4 to 43.7)
Central Europe, eastern Europe, and central Asia				
All causes	12.4% (9.0 to 16.2)	6.9% (0.72 to 13.8)	4.8% (3.8 to 6.0)	4.8% (3.2 to 6.4)
Diarrhoeal diseases	61.2% (31.8 to 83.2)	18.9% (10.3 to 27.1)	44.8% (-1.8 to 81.1)	15.5% (9.9 to 20.1)
Lower respiratory infections	43.6% (33.0 to 57.4)	25.0% (0.94 to 52.4)	15.3% (10.0 to 20.5)	17.6% (11.5 to 23.7)
Malaria	0.0% (0 to 0)	0.0% (0 to 0)	..	0.0% (0 to 0)
Measles	40.2% (19.5 to 49.8)	13.6% (6.4 to 18.0)	14.4% (5.5 to 26.4)	19.7% (9.0 to 26.4)
High income				
All causes	1.2% (0.82 to 1.9)	0.6% (0.2 to 1.4)	0.5% (0.4 to 0.7)	0.4% (0.3 to 0.5)
Diarrhoeal diseases	32.7% (12.1 to 60.1)	6.0% (3.0 to 9.0)	23.9% (-0.52 to 58.1)	6.4% (3.8 to 8.4)
Lower respiratory infections	34.8% (24.6 to 54.8)	17.7% (0.5 to 47.4)	8.8% (5.2 to 12.3)	13.7% (10.1 to 17.9)
Malaria	4.2% (-1.8 to 13.4)	3.8% (-1.5 to 11.7)	..	0.5% (-0.2 to 2.0)
Measles	14.2% (6.2 to 18.4)	4.0% (1.8 to 5.4)	5.5% (2.0 to 10.3)	5.5% (2.4 to 7.6)
Latin America and Caribbean				
All causes	10.3% (7.8 to 12.9)	6.4% (3.0 to 10.0)	4.4% (3.1 to 5.5)	4.0% (2.9 to 5.1)
Diarrhoeal diseases	56.8% (38.6 to 75.5)	20.7% (11.9 to 29.7)	32.6% (-1.1 to 69.0)	20.3% (13.7 to 26.0)
Lower respiratory infections	52.4% (38.9 to 68.9)	31.5% (1.3 to 63.4)	10.0% (6.0 to 13.6)	27.0% (19.0 to 33.7)
Malaria	14.4% (-7.1 to 45.4)	10.3% (-5.0 to 29.4)	..	5.6% (-2.4 to 24.8)
Measles	27.3% (12.9 to 34.5)	8.5% (4.0 to 11.2)	6.5% (2.5 to 12.3)	15.3% (6.9 to 21.6)
North Africa and Middle East				
All causes	10.9% (7.9 to 13.8)	6.9% (3.3 to 10.3)	5.7% (4.0 to 7.4)	4.5% (3.3 to 5.9)
Diarrhoeal diseases	68.4% (43.9 to 83.7)	29.6% (19.1 to 40.0)	47.8% (-2.1 to 81.2)	23.9% (16.4 to 29.7)
Lower respiratory infections	54.3% (44.1 to 63.4)	34.7% (2.0 to 58.9)	20.8% (13.9 to 27.1)	27.7% (19.2 to 35.0)
Malaria	31.3% (-22.7 to 78.3)	24.3% (-16.2 to 58.1)	..	13.1% (-6.7 to 50.9)
Measles	61.4% (34.5 to 71.2)	30.0% (15.2 to 38.3)	22.4% (9.3 to 38.9)	33.5% (15.6 to 41.5)
South Asia				
All causes	14.6% (11.8 to 16.8)	10.0% (4.3 to 13.2)	8.7% (6.3 to 10.4)	6.1% (4.4 to 7.9)
Diarrhoeal diseases	79.4% (58.3 to 88.3)	42.8% (29.6 to 54.0)	58.6% (-3.9 to 86.6)	27.7% (19.6 to 34.2)
Lower respiratory infections	53.4% (45.4 to 59.0)	38.4% (2.9 to 56.2)	26.9% (19.5 to 33.4)	27.0% (17.4 to 34.0)
Malaria	32.1% (-24.3 to 78.9)	25.0% (-17.4 to 60.1)	..	13.8% (-7.3 to 51.8)
Measles	75.9% (49.1 to 85.7)	42.5% (23.9 to 51.6)	36.0% (16.5 to 55.4)	38.4% (19.2 to 46.8)
Southeast Asia, east Asia, and Oceania				
All causes	12.3% (9.1 to 15.2)	7.7% (3.1 to 11.2)	5.9% (4.1 to 7.7)	5.2% (3.8 to 6.8)
Diarrhoeal diseases	73.0% (50.6 to 86.9)	32.5% (20.3 to 43.5)	49.1% (-2.2 to 84.1)	25.5% (17.3 to 31.7)
Lower respiratory infections	56.8% (43.9 to 66.3)	38.0% (2.2 to 62.9)	19.6% (13.0 to 25.3)	28.2% (19.6 to 35.1)
Malaria	25.7% (-17.2 to 68.9)	18.1% (-10.6 to 47.0)	..	12.3% (-6.1 to 46.3)
Measles	61.5% (35.8 to 70.9)	29.7% (15.8 to 37.2)	20.6% (8.6 to 35.3)	33.3% (16.4 to 41.2)

(Table 4 continues on next page)

children, could change our understanding of the current and forecasted burden of CGF. Third, although we believe that our analysis strengthens previous burden estimates by using estimated continuous distributions for CGF prevalence and for the risk of infections associated with CGF, our approach is cross-sectional in time and might not reflect the longitudinal nature of childhood growth. For example, a child who has been experiencing growth failure for several months might have a different risk of

infection or death than a child with a newly incident case of growth failure—a nuance that is not captured in our analysis. Such an analysis would need to account for repeated measurements of growth and potentially repeated measures of infection events.⁴³ It is likely that a cross-sectional measurement of growth does not accurately represent an individual child's risk of disease, especially given the importance of birth and early-life exposures.³⁵ Fourth, as mentioned earlier, low birthweight and short

	Child growth failure	Child underweight	Child wasting	Child stunting
(Continued from previous page)				
Sub-Saharan Africa				
All causes	23.4% (11.3 to 32.8)	15.1% (5.6 to 22.3)	10.7% (6.1 to 15.6)	10.0% (6.3 to 14.6)
Diarrhoeal diseases	77.3% (55.2 to 90.2)	37.0% (24.5 to 48.6)	53.1% (-3.0 to 87.0)	27.0% (19.4 to 33.3)
Lower respiratory infections	65.2% (52.5 to 73.9)	44.7% (2.9 to 69.7)	22.9% (15.1 to 30.1)	37.2% (30.6 to 43.2)
Malaria	26.3% (-17.7 to 69.4)	18.9% (-11.2 to 48.0)	..	12.4% (-6.3 to 45.8)
Measles	66.2% (39.5 to 75.8)	33.9% (18.1 to 42.2)	24.3% (10.4 to 40.2)	35.4% (17.4 to 43.7)

Estimates combine burden associated with mild, moderate, and severe forms of child growth failure: stunting was defined as height-for-age Z score (HAZ) less than -1; underweight was defined as weight-for-age Z score (WAZ) less than -1; and wasting was defined as weight-for-height Z score (WHZ) less than -1, according to WHO Child Growth Standards. The results presented herein incorporate between-study variation, meaning the uncertainty intervals here represent both uncertainty in the relative risks as well as predicted variation due to between-study variation, and as such some uncertainty intervals' lower bounds fall below zero. As described in the publication associated with the method,²⁴ a risk factor, outcome, or cause combination would not be included in the table if the uncertainty without between-study variation was not statistically significantly different from zero. Data in parentheses are 95% uncertainty intervals. Population attributable fractions are presented to one decimal place. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.

Table 4: All-cause and cause-specific population attributable fraction of deaths among children younger than 5 years for child growth failure at the global and GBD super-regional levels, 2023




Figure 3: Deaths from all child growth failure (mild, moderate, and severe) in children younger than 5 years per 100 000 in 2023
CGF=child growth failure.

gestation have been found to be correlated with CGF outcomes at birth, during the critical first 1000 days of life, and throughout childhood. Better data and a deeper analysis are needed to carefully link health status at and before birth to health outcomes throughout childhood to parse the relative risks of wasting, stunting, and underweight from low birthweight and short gestation. Finally, we used an adjustment after modelling our relative risk curves to

account for the correlation between different growth failure indicators. For example, children who have a low height-for-age are more likely to have low weight-for-age.¹² Future work to simultaneously quantify such correlations in a single statistical model might have an important impact on risk estimates.

The global burden of disease associated with all CGF (mild, moderate, and severe) is substantial and

concentrated in south Asia and sub-Saharan Africa. Children experiencing growth failure are at increased risk of mortality and incidence of infectious diseases. Although the burden associated with CGF has decreased in the past 20 years, more must be done to prevent children from being born small or preterm and to prevent children's growth from faltering. All children deserve an opportunity to have a healthy and productive life, but too many are being denied that chance because of poor growth. By accurately parsing the relative associations of the three main CGF indicators with childhood mortality and morbidity, our estimates provide an unprecedented opportunity to target interventions specifically for the dominant indicator, outcome, and cause of burden in each location.

GBD 2023 Child Growth Failure Collaborators

Christopher E Troeger*, Michael Benjamin Arndt*, Hasan Aalruz, Meriem Abdoun, Auwal Abdullahi, Mesfin Abebe, Armita Abedi, Alemwork Abie, Richard Gyan Aboagye, Hassan Abolhassani, Yonas Derso Abtew, Ahmed Abu-Zaid, Lawan Hassan Adamu, Mesafint Molla Adane, Isaac Yeboah Addo, Oyelola A Adegbeye, Victor Adekanmbi, Juliana Bunmi Adetunji, Qorinah Estiningtyas Sakilah Adnani, Leticia Akua Adzignbli, Muhammad Sohail Afzal, Saira Afzal, Navidha Aggarwal, Aqeel Ahmad, Muayyad M Ahmad, Sajjad Ahmad, Elham Ahmadi, Ayman Ahmed, Haroon Ahmed, Mehrunnisha Sharif Ahmed, Mushood Ahmed, Marjan Ajami, Budi Aji, Syed Mahfuz Al Hasan, Omar Al Omari, Mohammad Khursheed Alam, Mohammed Albashtawy, Fentahun Alemnew, Ayman Al-Eyadhy, Mohammed Usman Ali, Rafat Ali, Syed Shujait Ali, Waad Ali, Joseph Uy Almazan, Hesham M Al-Mekhlafi, Mohammed A Alsabri, Najim Z Alshahrani, Awais Altaf, Nelson Alvis-Guzman, Mohammad Al-Wardat, Hany Aly, Dickson A Amugsi, Abhishek Anil, Zelalem Alamrew Anteneh, Boluwatife Stephen Anuoluwa, Saeid Anvari, Anayochukwu Edward Anyasodor, Jalal Arabloo, Aleksandr Y Aravkin, Demelash Areda, Mahwish Arooj, Anton A Artamonov, Ashokan Arumugam, Nurila Aryntayeva, Bernard Kwadwo Yeboah Asiamah-Asare, Seyyed Shamsadin Athari, Maha Moh'd Wahbi Atout, Amlaku Mulat Aweke, Adedapo Wasiu Awotidebe, Asteray Assmie Ayenew, Priyamadhaba Behera, Michelle L Bell, Maryam Bemanalizadeh, Ajeet Singh Bhadoria, Sonu Bhaskar, Priyadarshini Bhattacharjee, Jasvinder Singh Bhatti, Catherine Bisignano, Bijit Biswas, Trupti Bodhare, Srinivasa Rao Bolla, Sri Harsha Boppana, Angelo Capodici, Rama Mohan Chandika, Vijay Kumar Chattu, Anis Ahmad Chaudhary, Moges Sisay Chekole, Hana Chen, Daniel Youngwhan Cho, Sonali Gajanan Choudhari, Isaac Sunday Chukwu, Erin Chung, Natalia Cruz-Martins, Alanna Gomes da Silva, Tukur Dahiru, Xiaochen Dai, Lalit Dandona, Rakhi Dandona, Samuel Demissie Darcho, Amira Hamed Darwish, Fernando Pio De la Hoz, Edgar Denova-Gutiérrez, Vinoth Gnana Chellaiyan Devanbu, Devananda Devegowda, Adriana Dima, Thanh Chi Do, Robert Kokou Dowou, Angel Belle Cheng Dy, Ibrahim Farahat El Bayoumy, Marwa Eldegwi, Mohammed Elhadi, Legesse Tesfaye Elilo, Iman El Sayed, Abidemi Omolara Fasanmi, Marta Figueiredo, Florian Fischer, Artem Alekseevich Fomenkov, Amin Fraij, Amanuel Tesfay Gebremedhin, Lemma Getacher, Genanew K Getahun, Maryam Gholamalizadeh, Nora M Gilbertson, Alem Abera Girmay, Mahaveer Golechha, Dinorah Gonzalez-Castell, Michal Grivna, Shi-Yang Guan, Mohammed Ibrahim Mohialdeen Gubari, Damitha Asanga Gunawardane, Zhifeng Guo, Bhawna Gupta, Rajat Das Gupta, Demewoz Haile, Nadia M Hamdy, Alexis J Handal,

Nasrin Hanifi, Habtamu Endashaw Hareru, Eka Mishbahatul Marah Has, Ahmed I Hasaballah, Ikrama Hassan, Simon I Hay, Khezar Hayat, Jiawei He, Behzad Heibati, Austin Heuer, Kamal Hezam, Ramesh Holla, Md Sabbir Hossain, Hassan Hosseinzadeh, Sorin Hostiuc, Tanvir M Huda, Javid Hussain, Dursa Hussein, Hong-Han Huynh, Bing-Fang Hwang, Segun Emmanuel Ibitoye, Mustapha Immurana, Teresa R Iskander, Md Rabiul Islam, Md Sahidul Islam, Sheikh Mohammed Shariful Islam, Louis Jacob, Mihajlo Jakovljevic, Shubha Jayaram, Achala Upendra Jayatilleke, Wenyi Jin, Alex Joseph, Nitin Joseph, Ali Kabir, Vidya Kadashetti, Dler H Hussein Kadir, Sanjay Kalra, Arun Kamireddy, Kehinde Kazeem Kanmodi, Rami S Kantar, Faizan Zaffar Kashoo, Gbenga A Kayode, Shemsu Kedir, Tibebelesassie S Keflie, Ajmal Khan, Maseer Khan, Vishnu Khanal, Shaghayegh Khanmohammadi, Khaled Khatab, Moawiah Mohammad Khatatbeh, Mahalaqua Nazli Khatib, Feriha Fatima Khidri, Kwanghyun Kim, Min Seo Kim, Adnan Kisa, Farzad Kompani, Isaac Koomson, Kewal Krishan, Mukhtar Kulimbet, Dewesh Kumar, G Anil Kumar, Nithin Kumar, Vijay Kumar, Almagul Kurmanova, Maria Dyah Kurniasari, Dian Kusuma, Chandrakant Lahariya, Kamaluddin Latief, Minh Huu Nhat Le, Nhi Huu Hanh Le, Sang-woong Lee, Seung Won Lee, Yo Han Lee, Virendra S Ligade, Stephen S Lim, Jialing Lin, Jue Liu, Xuefeng Liu, Rakesh Lodha, José Francisco López-Gil, Surbala Devi Lourembam, Zheng Feei Ma, Mahmoud Mabrok, Kashish Malhotra, Ahmad Azam Malik, Wahid Mansouri, Emmanuel Manu, Melvin Barrientos Marzan, Roy Rillera Marzo, Sammer Marzouk, Medha Mathur, Rita Mattiello, Rishi P Mediratta, Riffat Mehboob, Kala M Mehta, Tesfahun Mekene Meto, Tomislav Mestrovic, Sachith Mettananda, Tomasz Miazgowski, Giuseppe Minervini, Mojgan Mirghafourvand, Andreea Mirica, Jama Mohamed, Nouh Saad Mohamed, Sakineh Mohammad-Alizadeh-Charandabi, Abdollah Mohammadian-Hafshejani, Shafiu Mohammed, Ali H Mokdad, Lorenzo Monasta, Mohammad Ali Moni, Sumaira Mubarik, Sumoni Mukherjee, Sileshi Mulatu, Francesk Mulita, Christopher J L Murray, Ghulam Mustafa, Ayoub Nafei, Ganesh R Naik, Zuhair S Natto, Javid Nauman, Samidi Nirasha Kumari Navaratna, Biswa Prakash Nayak, Amanuel Tebabal Nega, Samata Nepal, Henok Biresaw Netsere, Georges Nguefack-Tsague, Dang Nguyen, The Phuong Nguyen, Robina Khan Niazi, Ali Nikoobar, Lawrence Achilles Nnyanzi, Shuhei Nomura, Mehran Nouri, Chisom Adaobi Nri-Ezedi, Dieta Nurrika, Sylvester Dodzi Nyadanu, Chimezie Igwegbe Nzopotam, Ogochukwu Janet Nzopotam, Ismail A Odetokun, Hassan Okati-Aliabad, Akinkunmi Paul Okekunle, Osarein Christabel Okonji, Bolajoko Olubukunola Olusanya, Jacob Olusegun Olusanya, Uchechukwu Levi Osuagwu, Amel Ouyahia, Mahesh P A, Jagadish Rao Padubidri, Anca Pantea Stoian, Romil R Parikh, Jay Patel, Shankargouda Patil, Shrikant Pawar, Gavin Pereira, Arokiasamy Perianayagam, Fanny Emily Petermann-Rocha, Hoang Nhat Pham, Hoang Tran Pham, My Kieu Phan, Jalandhar Pradhan, Pranil Man Singh Pradhan, Akila Prashant, Jagadeesh Puvvula, Ibrahim Qattea, Pankaja Raghav, Md. Mosfequr Rahman, Mosiur Rahman, Muhammad Aziz Rahman, Amir Masoud Rahmani, Masoud Rahmati, Rajesh Kumar Rai, Ivano Raimondo, Sathish Rajaa, Rayan Rajabi, Mahmoud Mohammed Ramadan, Chitra Ramasamy, Shakthi Kumaran Ramasamy, Chhabi Lal Ranabhat, Chythra R Rao, Sowmya J Rao, Davide Rasella, Mamunur Rashid, Devarajan Rathish, Santosh Kumar Rauniyar, David Laith Rawaf, Salman Rawaf, Nazila Rezaei, Mohsen Rezaeian, Thales Philippe Rodrigues da Silva, Jefferson Antonio Buendia Rodriguez, Peter Rohloff, Debby Syahrul Romadlon, Bedanta Roy, Shubhanjali Roy, Cameron John Sabet, Kabir P Sadarangani, Basema Ahmad Saddik, Umar Saeed, Amene Saghazadeh, Dominic Sagoe, Narjes Saheb Sharif-Askari, Amirhossein Sahebkar, Pragnay Monalisa Sahoo, Yoseph Leonardo Samodra, Abdallah M Samy, Rama Krishna Sanjeev, Senthilkumar Sankararaman, Milena M Santric-Milicevic, Jacob Owusu Sarfo, Yaser Sarikhani, Tanmay Sarkar, Gargi Sachin Sarode, Sachin C Sarode, Benn Sartorius, Jennifer Saulam, Monika Sawhney, Ganesh Kumar Saya, Christophe Schinckus, Art Schuermans, Ashenafi Kibret Sendekie,

Subramanian Senthilkumaran, Yashendra Sethi, Allen Seylani, Shazlin Shaharudin, Samiah Shahid, Masood Ali Shaikh, Sunder Sham, Muhammad Aaqib Shamim, Mohd Shanawaz, Mohammed Shannawaz, Nigussie Tadesse Sharew, Vishal Sharma, Pavanchand H Shetty, Aminu Shittu, Ivy Shiue, Seyed Afshin Shorofi, Emmanuel Edwar Siddig, Mithun Sikdar, Luís Manuel Lopes Rodrigues Silva, Harmanjit Singh, Jasvinder A Singh, Kalpana Singh, Surjit Singh, Shipra Solanki, Mansi Soni, Reed J D Sorensen, Muhammad Suleman, Desy Sulistiyorini, Chandan Kumar Swain, Seyyed Mohammad Tabatabaei, Seyed-Amir Tabatabaeizadeh, Mohammad Tabish, Jacques Lukenze Tamuzi, Birhan Tsegaw Taye, Wegayehu Zeneb Teklehaimanot, Abainash Tekola, Mohamad-Hani Temsah, Rekha Thapar, Jansje Henny Vera Ticoalu, Tenaw Yimer Tiruye, Mariya Vladimirovna Titova, Sojit Tomo, Marcos Roberto Tovani-Palone, Quynh Thuy Huong Tran, Thang Huu Tran, Nguyen Tran Minh Duc, Aristidis Tsatsakis, Abdul Rohim Tualeka, Saeed Ullah, Shahid Ullah, Muhammad Umair, Bhaskaran Unnikrishnan, Era Upadhyay, Jibrin Sammani Usman, Jef Van den Eynde, Siavash Vaziri, Balachandar Vellingiri, Vasily Vlassov, Gebeyaw Biset Wagaw, Yanzhong Wang, Felicia Wu, Hong Xiao, Vikas Yadav, Galal Yahya, Dong Keon Yon, Naohiro Yonemoto, Chuanhua Yu, Sojib Bin Zaman, Iman Zare, Michael Zastrozhin, Mohammed G M Zeariya, Salih M Mustafa Salih Zebari, Claire Chenwen Zhong, Nicholas J Kassebaum, Robert C Reiner Jr**
*First authors. **Senior author.

Affiliations

See Online for appendix 2
Please see appendix 2 (pp 4–16) for the affiliations for individual authors.

Contributors

Please see appendix 2 (pp 17–24) for more detailed information about individual author contributions to the research, divided into the following categories: managing the overall research enterprise; writing the first draft of the manuscript; primary responsibility for applying analytical methods to produce estimates; primary responsibility for seeking, cataloguing, extracting, or cleaning data; designing or coding figures and tables; providing data or critical feedback on data sources; developing methods or computational machinery; providing critical feedback on methods or results; drafting the manuscript or revising it critically for important intellectual content; and managing the estimation or publications process. The lead and senior authors had full access to the data in the study and final responsibility for the decision to submit the manuscript for publication.

Declaration of interests

S Afzal reports support for the present manuscript from the Institute of Public Health Lahore; grants or contracts from the Dean office Institute of Public Health Lahore; honoraria for experts, lectures, visiting speakers, and educational seminars provided by the Dean Institute of Public Health Lahore; support for attending meetings and travel provided by the Dean Institute of Public Health, Lahore Pakistan; leadership or fiduciary roles in board, society, committee, or advocacy groups, paid or unpaid as a Member of the Pakistan Higher Education Commission Research Committee, Member of Pakistan Medical and Dental Commission Research and Journals Committee, Member of Pakistan National Bioethics Committee, Member of Pakistan Society of Internal Medicine, Member of Pakistan Association of Medical Editors, Member of Medical Microbiology and Infectious Diseases Society, a Fellow of LEADS International, Fellow of Faculty of Public Health UK, and as a Fellow of College of Physicians and Surgeons Pakistan; receipt of equipment, materials, drugs, and services including computer software and equipment from Bergen University Norway for research writing; and other financial or non-financial support from the Dean Public Health Institute of Public Health Birdwood Lahore, outside the submitted work. J Baker reports grants or contracts paid to her institution from Novo Nordisk Foundation, World Cancer Research Fund, Independent Research Council Denmark, and European Union Horizon; consulting fees from Novo Nordisk; payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing, or educational events from Novo Nordisk; support for attending meetings or travel, or both, from the European Association for the Study of Obesity; participation on a data safety monitoring board or advisory

board from Novo Nordisk; and leadership or fiduciary roles in other board, society, committee, or advocacy groups, paid or unpaid with the European Association for the Study of Obesity, outside the submitted work. M L Bell reports grants or contracts paid to her institution from the US Environmental Protection Agency, National Institutes of Health, Hutchinson Postdoctoral Fellowship, Health Effects Institute, Robert Wood Johnson Foundation, Wellcome Trust Foundation, and Internal Yale funding for the Yale Institute for Biospheric Studies; consulting fees from Clinique and Toximap; honoraria for speaking from the Colorado School of Public Health, Duke University, University of Texas, Data4Justice, Korea University, UPenn, Brown University, and Northeastern University; honorarium for editorial duties from IOP Publishing; honorarium for grant review from the NIH, Health Canada, Environment, Health & Safety, Program Advisory Committee, UK Research and Innovation, AXA Research Fund Fellowship, and University of Texas; honorarium for research from Korea University; honoraria for participation on an external advisory committee from Harvard University and University of Montana; honorarium for online survey or workshop from SciQuest; support for attending meetings or travel, or both, from the Colorado School of Public Health, University of Texas, Duke University, Harvard University, American Journal of Public Health, Columbia University, Harvard, Community Modeling and Analysis System conference, Nature Conference, Boston University, and Northeastern University; leadership or fiduciary roles in other board, society, committee, or advocacy groups, paid or unpaid from the Fifth National Climate Assessment, *Lancet* Countdown, US EPA Clean Air Scientific Advisory Committee (CASAC), Johns Hopkins Department of Environmental Health And Engineering Advisory Board, Harvard external advisory committee for training grant, WHO Global Air Pollution and Health Technical Advisory group, and the National Academies Panels and Committees, outside the submitted work. S Bhaskar reports grants or contracts from the Japan Society for the Promotion of Science (JSPS), Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT), Grant-in-Aid for Scientific Research (KAKENHI; Grant ID 23KF0126), JSPS and the Australian Academy of Science, JSPS International Fellowship (Grant ID P23712); leadership or fiduciary roles in other board, society, committee, or advocacy groups, paid or unpaid as District Chair, Diversity, Equity, Inclusion & Belonging of Rotary District 9675 (Sydney, Australia); as Chair, Founding Member, and Manager of the Global Health & Migration Hub Community, Global Health Hub Germany (Berlin, Germany); as Editorial Board Member of *PLOS One*, *BMC Neurology*, *Frontiers in Neurology*, *Frontiers in Stroke*, *Frontiers in Public Health*, *Journal of Aging Research*, *Neurology International*, *Diagnostics*, and *BMC Medical Research Methodology*; as a member of the College of Reviewers, Canadian Institutes of Health Research (CIHR), Government of Canada; as the Director of Research of World Headache Society (Bengaluru, India), as Expert Adviser/Reviewer of Cariplo Foundation (Milan, Italy), as Visiting Director of National Cerebral and Cardiovascular Center, Department of Neurology, Division of Cerebrovascular Medicine and Neurology, Suita (Osaka, Japan); as Member, Scientific Review Committee of Cardiff University Biobank (Cardiff, UK), as Chair of Rotary Reconciliation Action Plan, and Healthcare and Medical Adviser at Japan Connect (Osaka, Japan), outside the submitted work. A A Fomenkov reports support for the present manuscript from the Ministry of Science and Higher Education of the Russian Federation (theme number 122042600086-7 and number 122042700043-9). N J Kassebaum reports support for the present manuscript from the Gates Foundation (funding for Global Burden of Disease Study); consulting fees from Fujifilm Sonosite (product development consultation unrelated to present work) and Philips (product development consultation unrelated to present work); payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing, or educational events from Bristol Myers Squibb (travel expenses plus consulting fees for presenting GBD results at international meeting). K Krishan reports other financial or non-financial interests from the UGC Centre of Advanced Study, CAS II, awarded to the Department of Anthropology, Panjab University (Chandigarh, India), outside the submitted work. J Liu reports support for the present manuscript and grants from the National Natural Science Foundation (72474005) and Beijing Natural Science Foundation

(L222027). L Monasta reports support for the present manuscript from the Italian Ministry of Health (Ricerca Corrente 34/2017), payments made to the Institute for Maternal and Child Health IRCCS Burlo Garofolo. S Nomura reports support for the present manuscript from the Ministry of Education, Culture, Sports, Science and Technology of Japan (24H00663) Grant and the Precursory Research for Embryonic Science and Technology from the Japan Science and Technology Agency (JPMJPR22R8) Grant. Y L Samodra reports grants or contracts from Taipei Medical University (Taiwan), Type A Doctoral Scholarship, Institute of Epidemiology and Preventive Medicine, National Taiwan University, National Science and Technology Council Taiwan, Post-doctoral Fellow contract; payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing, or educational events from Generasi Peneliti, Indonesia; leadership or fiduciary roles in board, society, committee, or advocacy groups, paid or unpaid as the co-founder of Benang Merah Research Center, Indonesia (<https://www.benangmerah.net>); and other financial or non-financial interests as a scholarship mentor with Jago Beasiswa (<https://www.idebeasiswa.com>), outside the submitted work. V Sharma acknowledges support from Directorate of Forensic Science, Ministry of Home Affairs research project (DFSS28(1)2019/EMR/6) at the Institute of Forensic Science & Criminology, Panjab University (Chandigarh, India), outside the submitted work. L M L R da Silva reports grants or contracts from SPRINT (Sport Physical Activity and Health Research & Innovation Center), Polytechnic of Guarda (6300-559 Guarda, Portugal) and RISE-Health, Faculty of Health Sciences, University of Beira Interior (6201-506 Covilhã, Portugal), outside the submitted work. J Singh reports consulting fees from ROMTech, Atheneum, Clearview Healthcare Partners, American College of Rheumatology, Yale, Hulio, Horizon Pharmaceuticals, DINORA, ANI/Exeltis, USA Inc, Frictionless Solutions, Schipfer, Crealta/Horizon, Medisys, Fidia, PK Med, Two labs, Adept Field Solutions, Clinical Care Options, Putnam Associates, Focus Forward, Navigant Consulting, Spherix, MediQ, Jupiter Life Science, UBM LLC, Trio Health, Medscape, WebMD, and Practice Point Communications; and the NIH; payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing, or educational events from Simply Speaking; support for attending meetings or travel, or both, from Simply Speaking; leadership or fiduciary roles in other board, society, committee, or advocacy groups, paid or unpaid as a past steering committee member of OMERACT, an international organization that develops measures for clinical trials and receives arm's length funding from 12 pharmaceutical companies, and as Chair of the Veterans Affairs Rheumatology Field Advisory Committee, and as editor and Director of the UAB Cochrane Musculoskeletal Group Satellite Center on Network Meta-analysis; stock or stock options in Atai Life Sciences, Kintara Therapeutics, Intelligent Biosolutions, Acumen Pharmaceutical, TPT Global Tech, Vaxart Pharmaceuticals, Atyu Biopharma, Adaptimmune Therapeutics, GeoVax Labs, Pieris Pharmaceuticals, Enzolytics, Seres Therapeutics, Tonix Pharmaceuticals Holding Corp, Aebona Pharmaceuticals, and Charlotte's Web Holdings; and previously owned stock options in Amarin, Viking, and Moderna Pharmaceuticals, outside the submitted work. J H V Ticoalu reports leadership or fiduciary roles in board, society, committee, or advocacy groups, paid or unpaid as Co-founder of Benang Merah Research Center, Indonesia (<https://www.benangmerah.net>), outside the submitted work. M Titova reports support for the present manuscript from the state assignment of the Ministry of Science and Higher Education of the Russian Federation (theme number 122042600086-7 and number 122042700043-9). E Upadhyay reports the following patents planned, issued or pending with The Office of the Controller General of Patents, Designs & Trade Marks (CGPDTM; <https://iprsearch.ipindia.gov.in/PublicSearch/PublicationSearch/ApplicationStatus>): "Eco-friendly bio-shoe polish from banana and turmeric" (filed 202511021382), "Honey-based polyherbal syrup composition to treat air pollution-induced inflammation and preparation method thereof" (filed 202511035171), "Process for preparing a caffeine free, antioxidant and nutrient rich beverage" (filed 202511042794), "A system and method of reusable filters for anti-pollution mask" (published 202011003559), "A system and method for electricity generation through crop stubble by using microbial fuel cells" (published 202011008531), "A system for disposed personal protection equipment (PPE) into biofuel through pyrolysis and

method" (published 202111005659), "A novel herbal pharmaceutical aid for formulation of gel and method thereof" (published 202111023333), "Herbal drug formulation for treating lung tissue degenerated by particulate matter exposure" (published 202311035276), "A method to transform cow dung into the wall paint by using natural materials and composition thereof" (filed 202311085452), "Biodegradable packaging composition and method of preparation thereof" (filed 202511017848); and leadership or fiduciary roles in other board, society, committee, or advocacy groups as an Executive Council Member for the Indian Meteorological Society, Jaipur Chapter (India) and a Member Secretary for the Department of Science & Technology (DST), Promotion of University Research and Scientific Excellence (PURSE) Program, outside the submitted work.

Data sharing

To download the citations and metadata for the input data used in these analyses, please visit the Global Health Data Exchange GBD 2023 Sources Tool: <https://ghdx.healthdata.org/gbd-2023/sources>. To download estimates produced in these analyses, please visit the GBD Results Tool: <https://vizhub.healthdata.org/gbd-results/>.

Acknowledgments

The research reported in this publication was supported by the Gates Foundation (OPP1152504).

References

- Bhutta ZA, Berkley JA, Bandsma RHJ, Kerac M, Trehan I, Briend A. Severe childhood malnutrition. *Nat Rev Dis Primers* 2017; 3: 17067.
- GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; 396: 1223–49.
- Olofin I, McDonald CM, Ezzati M, et al, and the Nutrition Impact Model Study (anthropometry cohort pooling). Associations of suboptimal growth with all-cause and cause-specific mortality in children under five years: a pooled analysis of ten prospective studies. *PLoS One* 2013; 8: e64636.
- Black RE, Victora CG, Walker SP, et al, and the Maternal and Child Nutrition Study Group. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* 2013; 382: 427–51.
- McDonald CM, Olofin I, Flaxman S, et al, and the Nutrition Impact Model Study. The effect of multiple anthropometric deficits on child mortality: meta-analysis of individual data in 10 prospective studies from developing countries. *Am J Clin Nutr* 2013; 97: 896–901.
- Victora CG, Adair L, Fall C, et al, and the Maternal and Child Undernutrition Study Group. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* 2008; 371: 340–57.
- GBD 2019 Under-5 Mortality Collaborators. Global, regional, and national progress towards Sustainable Development Goal 3.2 for neonatal and child health: all-cause and cause-specific mortality findings from the Global Burden of Disease Study 2019. *Lancet* 2021; 398: 870–905.
- WHO. Fourth round of the global pulse survey on continuity of essential health services during the COVID-19 pandemic: November 2022–January 2023. May 1, 2023. https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS_continuity-survey-2023.1 (accessed Jan 15, 2025).
- Reiner RC Jr, Hay SI, Hay SI, and the LBD Triple Burden Collaborators. The overlapping burden of the three leading causes of disability and death in sub-Saharan African children. *Nat Commun* 2022; 13: 7457.
- Kramer MS, Kakuma R. Optimal duration of exclusive breastfeeding. *Cochrane Database Syst Rev* 2012; 2012: CD003517.
- Soofi SB, Khan GN, Ariff S, et al. Effectiveness of nutritional supplementation during the first 1000-days of life to reduce child undernutrition: a cluster randomized controlled trial in Pakistan. *Lancet Reg Health Southeast Asia* 2022; 4: 100035.
- Mertens A, Benjamin-Chung J, Colford JM Jr, et al, and the Ki Child Growth Consortium. Child wasting and concurrent stunting in low- and middle-income countries. *Nature* 2023; 621: 558–67.

- 13 WHO. WHO guideline on the prevention and management of wasting and nutritional oedema (acute malnutrition) in infants and children under 5 years. Dec 29, 2023. <https://www.who.int/publications/i/item/9789240082830> (accessed Jan 14, 2025).
- 14 Suárez-Idueta L, Yargawa J, Blencowe H, et al, and the National Vulnerable Newborn Prevalence Collaborative Group and Vulnerable Newborn Measurement Core Group. Vulnerable newborn types: analysis of population-based registries for 165 million births in 23 countries, 2000–2021. *BJOG* 2023; published online May 8. <https://doi.org/10.1111/1471-0528.17505>.
- 15 Lawn JE, Ohuma EO, Bradley E, et al, the Lancet Small Vulnerable Newborn Steering Committee, and WHO/UNICEF Preterm Birth Estimates Group, and National Vulnerable Newborn Measurement Group, and the Subnational Vulnerable Newborn Measurement Group. Small babies, big risks: global estimates of prevalence and mortality for vulnerable newborns to accelerate change and improve counting. *Lancet* 2023; **401**: 1707–19.
- 16 Young MF, Nguyen PH, Gonzalez Casanova I, et al. Role of maternal preconception nutrition on offspring growth and risk of stunting across the first 1000 days in Vietnam: a prospective cohort study. *PLoS One* 2018; **13**: e0203201.
- 17 Petri WA Jr, Miller M, Binder HJ, Levine MM, Dillingham R, Guerrant RL. Enteric infections, diarrhea, and their impact on function and development. *J Clin Invest* 2008; **118**: 1277–90.
- 18 Kinyoki DK, Osgood-Zimmerman AE, Pickering BV, et al, and the Local Burden of Disease Child Growth Failure Collaborators. Mapping child growth failure across low- and middle-income countries. *Nature* 2020; **577**: 231–34.
- 19 Heidkamp RA, Piwoz E, Gillespie S, et al. Mobilising evidence, data, and resources to achieve global maternal and child undernutrition targets and the Sustainable Development Goals: an agenda for action. *Lancet* 2021; **397**: 1400–18.
- 20 Fitzgerald R, Manguerra H, Arndt MB, et al. Current dichotomous metrics obscure trends in severe and extreme child growth failure. *Sci Adv* 2022; **8**: eabm8954.
- 21 IHME. Protocol for the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD). March 2020. https://www.healthdata.org/sites/default/files/files/Projects/GBD/March2020_GBD%20Protocol_v4.pdf (accessed Aug 28, 2024).
- 22 GBD 2023 Disease and Injury and Risk Factor Collaborators. Burden of 375 diseases and injuries, risk-attributable burden of 88 risk factors, and healthy life expectancy in 204 countries and territories, including 660 subnational locations, 1990–2023: a systematic analysis for the Global Burden of Disease Study 2023. *Lancet* 2025; **406**: 1873–922.
- 23 GBD 2021 Risk Factors Collaborators. Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet* 2024; **403**: 2162–203.
- 24 Zheng P, Afshin A, Biryukov S, et al. The Burden of Proof studies: assessing the evidence of risk. *Nat Med* 2022; **28**: 2038–44.
- 25 GBD 2023 Causes of Death Collaborators. Global burden of 292 causes of death in 204 countries and territories and 660 subnational locations, 1990–2023: a systematic analysis for the Global Burden of Disease Study 2023. *Lancet* 2025; **406**: 1811–72.
- 26 GBD 2023 Demographics Collaborators. Global age-sex-specific all-cause mortality and life expectancy estimates for 204 countries and territories and 660 subnational locations, 1950–2023: a demographic analysis for the Global Burden of Disease Study 2023. *Lancet* 2025; **406**: 1731–810.
- 27 GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; **396**: 1204–22.
- 28 Weiss DJ, Lucas TCD, Nguyen M, et al. Mapping the global prevalence, incidence, and mortality of *Plasmodium falciparum*, 2000–17: a spatial and temporal modelling study. *Lancet* 2019; **394**: 322–31.
- 29 Stevens GA, Alkema L, Black RE, et al, and the GATHER Working Group. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet* 2016; **388**: e19–23.
- 30 GBD 2017 Diarrhoeal Disease Collaborators. Quantifying risks and interventions that have affected the burden of diarrhoea among children younger than 5 years: an analysis of the Global Burden of Disease Study 2017. *Lancet Infect Dis* 2020; **20**: 37–59.
- 31 GBD 2017 Lower Respiratory Infections Collaborators. Quantifying risks and interventions that have affected the burden of lower respiratory infections among children younger than 5 years: an analysis for the Global Burden of Disease Study 2017. *Lancet Infect Dis* 2020; **20**: 60–79.
- 32 Keats EC, Das JK, Salam RA, et al. Effective interventions to address maternal and child malnutrition: an update of the evidence. *Lancet Child Adolesc Health* 2021; **5**: 367–84.
- 33 Bhutta ZA, Akseer N, Keats EC, et al. How countries can reduce child stunting at scale: lessons from exemplar countries. *Am J Clin Nutr* 2020; **112** (suppl 2): 894S–904S.
- 34 Bhutta ZA, Ahmed T, Black RE, et al, and the Maternal and Child Undernutrition Study Group. What works? Interventions for maternal and child undernutrition and survival. *Lancet* 2008; **371**: 417–40.
- 35 Benjamin-Chung J, Mertens A, Colford JM Jr, et al, and the Ki Child Growth Consortium. Early-childhood linear growth faltering in low- and middle-income countries. *Nature* 2023; **621**: 550–57.
- 36 Georgiadis A, Penny ME. Child undernutrition: opportunities beyond the first 1000 days. *Lancet Public Health* 2017; **2**: e399.
- 37 Walson JL, Berkley JA. The impact of malnutrition on childhood infections. *Curr Opin Infect Dis* 2018; **31**: 231–36.
- 38 Kirolos A, Blacow RM, Parajuli A, et al. The impact of childhood malnutrition on mortality from pneumonia: a systematic review and network meta-analysis. *BMJ Glob Health* 2021; **6**: e007411.
- 39 de Onis M, Blössner M, Borghi E, Morris R, Frongillo EA. Methodology for estimating regional and global trends of child malnutrition. *Int J Epidemiol* 2004; **33**: 1260–70.
- 40 Barros AJD, Victora CG. Measuring coverage in MNCH: determining and interpreting inequalities in coverage of maternal, newborn, and child health interventions. *PLoS Med* 2013; **10**: e1001390.
- 41 Osendarp S, Akuoku JK, Black RE, et al. The COVID-19 crisis will exacerbate maternal and child undernutrition and child mortality in low- and middle-income countries. *Nat Food* 2021; **2**: 476–84.
- 42 Wang H, Paulson KR, Pease SA, et al, and the COVID-19 Excess Mortality Collaborators. Estimating excess mortality due to the COVID-19 pandemic: a systematic analysis of COVID-19-related mortality, 2020–21. *Lancet* 2022; **399**: 1513–36.
- 43 Anderson C, Hafen R, Sofrygin O, Ryan L, and the members of the HBGDki Community. Comparing predictive abilities of longitudinal child growth models. *Stat Med* 2019; **38**: 3555–70.