

# Hypoxia responsive gene expression is mediated by various subsets of transcription factors and miRNAs that are determined by the actual oxygen availability

Francesco Licausi<sup>1</sup>, Daan A. Weits<sup>1</sup>, Bikram Datt Pant<sup>1</sup>, Wolf-Rüdiger Scheible<sup>1</sup>, Peter Geigenberger<sup>2</sup> and Joost T. van Dongen<sup>1</sup>

<sup>1</sup>Max Planck Institute of Molecular Plant Physiology, Energy Metabolism & Molecular Genomics Research Groups, Am Mühlenberg 1, D-14476 Potsdam-Golm, Germany; <sup>2</sup>Department Biologie I, Ludwig-Maximilians-Universität München, D-82152 Martinsried, Germany

## Summary

Author for correspondence:

Francesco Licausi

Tel: +31 331 5678353

Email: [licausi@mpimp-golm.mpg.de](mailto:licausi@mpimp-golm.mpg.de)

Received: 11 June 2010

Accepted: 22 July 2010

*New Phytologist* (2011) **190**: 442–456

doi: 10.1111/j.1469-8137.2010.03451.x

**Key words:** anoxia, *Arabidopsis*, hypoxia, microRNA (miRNA), quantitative real-time PCR (qRT-PCR), transcription factor.

- Reduced oxygen availability is not only associated with flooding, but occurs also during growth and development. It is largely unknown how hypoxia is perceived and what signaling cascade is involved in activating adaptive responses.
- We analysed the expression of over 1900 transcription factors (TFs) and 180 microRNA primary transcripts (pri-miRNAs) in *Arabidopsis* roots exposed to different hypoxic conditions by means of quantitative PCR. We also analysed the promoters of genes induced by hypoxia with respect to over-represented DNA elements that can act as potential TF binding sites and their *in vivo* interaction was verified.
- We identified various subsets of TFs that responded differentially through time and in an oxygen concentration-dependent manner. The regulatory potential of selected TFs and their predicted DNA binding elements was validated. Although the expression of pri-miRNAs was differentially regulated under hypoxia, only one corresponding mature miRNA changed accordingly. Putative target transcripts of the miRNAs were not significantly affected.
- Our results show that the regulation of hypoxia-induced genes is controlled via simultaneous interaction of various combinations of TFs. Under anoxic conditions, an additional set of TFs is induced. Regulation of gene expression via miRNAs appears to play a minor role during hypoxia.

## Introduction

As ultimate electron acceptor of the oxidative phosphorylation reactions in mitochondria, oxygen represents an indispensable element for aerobic organisms, including plants. However, within plants, oxygen availability can vary strongly between tissues because of diffusional restrictions (Geigenberger, 2003, van Dongen *et al.*, 2003). Also, during organ development, the internal oxygen concentration can change considerably (Vigeolas *et al.*, 2003; Van Dongen *et al.*, 2004; Borisjuk *et al.*, 2007). Therefore, hypoxic conditions are common within plants even when oxygen is easily available from the environment. Moreover, severe hypoxic conditions or even anoxia can occur in plant tissues when the availability of environmental oxygen decreases, for example owing to reduced diffusion rates of gases in water during flooding or waterlogging (Armstrong, 1979).

Hypoxia not only represents a stress condition, but is also implicated as signal in the regulation of growth and development under optimal conditions. In human embryonic stem cells, for example, variations in oxygen levels are likely to play a role, by regulating pluripotency and proliferation (Forristal *et al.*, 2010). However, in plants, different cell types exhibit a conserved response to low oxygen levels at the molecular level (Mustroph *et al.*, 2010). This response includes the induction of a core set of genes after 30 min under hypoxia, whose expression is maintained for several hours (Klok *et al.*, 2002; van Dongen *et al.*, 2009). The increased transcript levels of these genes are further correlated with an enhanced association of the mRNAs with polysomes, suggesting that these transcripts are actively translated (Branco-Price *et al.*, 2008). Many of these genes that increase their expression when the oxygen availability reduces are required to maintain energy production via

glycolysis, such as pyruvate decarboxylase (*PDC*), alcohol dehydrogenase (*ADH*; Ismond *et al.*, 2003) and alanine aminotransferase (*AlaAT*; Rocha *et al.*, 2010).

As for many other responses to environmental stimuli, a signaling cascade is likely to be required to transmit the hypoxic signal from its perception to the core anaerobic response and integrate it in the developmental program. As the mechanism by which oxygen is perceived in plants is still largely unknown (Licausi & Perata, 2009), hypoxia-responsive transcription factors (TFs) are a good starting point to investigate the regulation of the hypoxic response, as they represent the last regulatory element in the signal cascade. Furthermore, analysis of *cis*-acting DNA sequences over-represented in anaerobic gene promoters can be coupled to the isolation of *trans*-acting TFs, to identify master regulators in the transcriptional cascade.

Various studies have addressed the role of some hypoxia responsive TFs belonging to different protein classes, including the MYB (Myeloblastosis), NAC [NAM (No Apical Meristem), ATAF (Arabidopsis Transcription Activation Factor), CUC (Cup-shaped Cotyledons)], PHD (Plant Homeodomain) and ERF (Ethylene Responsive Factor) families (Hoeren *et al.*, 1998; Bond *et al.*, 2009; Christianson *et al.*, 2009; Licausi *et al.*, 2010). However, to date, no TF has been shown to act alone to trigger the induction of the core-set of anaerobically induced genes. It therefore seems likely that a complex of transcriptional regulators is required to activate the molecular response to hypoxia. To understand the mechanisms and regulation of the molecular response to low oxygen, it is necessary to identify single members that comprise the TF complex first. Generally, TFs have very low transcript abundances, making proper detection a challenge. Although some changes in TF expression were identified via microarray approaches in the past, quantitative real-time PCR (qRT-PCR) is generally recognized to be better suited to identify differentially regulated TFs because of its higher sensitivity and accuracy (Czechowski *et al.*, 2002; Caldana *et al.*, 2007).

As TFs regulate the expression of other genes, including themselves or other TFs, reconstructing the hierarchy of their regulation can be a difficult task. Moreover, TFs constitute one of the widest functional classes of targets for small RNAs (Rhoades *et al.*, 2002). These noncoding transcripts are 20- to 24-nucleotide single-stranded RNAs, such as microRNAs (miRNAs) and *trans*-acting small interfering RNAs (ta-siRNAs) that promote degradation of the mRNAs to which they bind or hinder translation (Brodersen *et al.*, 2008; Lanet *et al.*, 2009). In plants, miRNAs have been shown to regulate development (Aukerman & Sakai, 2003; Emery *et al.*, 2003; Palatnik *et al.*, 2003) or the response to abiotic stresses (Sunkar & Zhu, 2004) including macronutrient limitation (Bari *et al.*, 2006; Chiou *et al.*, 2006; Pant *et al.*, 2008, 2009; Branscheid *et al.*, 2010; Huang *et al.*, 2010). Hypoxia-responsive miRNAs and ta-siRNAs

were reported by Moldovan *et al.* (2009). These noncoding RNAs might also account for the variations in mRNA levels of the anaerobic genes, especially TFs.

In the present study, we report the comprehensive analysis and identification of upregulated and downregulated transcripts for TFs and pri-miRNAs in roots of *Arabidopsis* seedlings treated for 30 min and 2 h with different hypoxic conditions using qRT-PCR. An extensive correlation analysis was performed on the TF/miRNA data and available microarray analyses obtained from identically treated plant material. Several conserved DNA-binding elements were identified and subsequently validated to be involved in the regulation of the transcriptional response of *Arabidopsis* roots to low oxygen.

## Materials and Methods

### Plant material and growth conditions

*Arabidopsis thaliana* ecotype Col-0 seedlings were grown in a similar manner to that described in van Dongen *et al.* (2009), with a 12 h light (160  $\mu$ E) and 12 h darkness cycle (day and night temperature was 22°C) on vertical plates with 10% agar dissolved in half-strength Murashige and Skoog (MS) medium (Duchefa Biochemie BV, Haarlem, the Netherlands) supplemented with 1% sucrose. The roots of the plants were growing on top of the agar, thus avoiding restricted oxygen diffusion towards the roots by the agar medium. Fifteen days after germination, seedlings were transferred to continuous darkness and a stream of pre-mixed air containing different oxygen concentrations (as indicated in legends to Figs 3 and 5), 350 ppm CO<sub>2</sub>, and N<sub>2</sub> (Air Liquide, Berlin, Germany) was blown over the plates. Each hypoxic treatment and the normoxic control treatments for all different time points were conducted similarly in darkness to abstract variation induced by variable factors other than the oxygen concentration, such as developmental variation or changes induced by the shift from light to dark. Roots were harvested in liquid nitrogen at the time-point indicated in the legends to Figs 3 and 5.

### RNA extraction and cDNA synthesis

Total RNA was isolated from roots using TRIZOL reagent (Invitrogen), as described by Czechowski *et al.* (2002). Seventy-five micrograms of total RNA were subjected to a DNase treatment using a TURBO DNA free kit (Ambion), according to the manufacturer's instructions. Absence of genomic DNA contamination was tested by PCR, using primers designed on intron sequence of a control gene (At5g65080). cDNA was synthesized from 13  $\mu$ g DNA-free RNA using SuperScript III reverse transcriptase (Invitrogen), according to the manufacturer's instructions. The efficiency of cDNA synthesis was assessed by real-time

PCR amplification of control genes encoding *UBQ10* (*At4g05320*) and 1 kb distant regions on the *GAPDH* (*At1g13440*) gene. Only cDNA preparations that yielded similar  $C_T$  (cycle number at which the fluorescence signal crosses a fixed threshold of 0.2 units) values ( $\pm 3$  cycles) for the *GAPDH* genes were used.

#### qRT-PCR for TF and pri-miRNA transcript quantification

Polymerase chain reactions were performed in an optical 384-well plate with an ABI PRISM 7900 HT Sequence Detection System (Applied Biosystems, Foster City, CA, USA), using SYBR Green to monitor dsDNA synthesis. Reactions contained 2.5  $\mu$ l 2 $\times$  FAST SYBR Green Master Mix reagent (Applied Biosystems), 13.0 ng cDNA and 100 nmol of each gene-specific primer in a final volume of 5  $\mu$ l. The following standard thermal profile was used for all PCRs: 95°C for 2 min; 40 cycles of 95°C for 3 s and 60°C for 20 s. In order to compare data from different PCR runs or cDNA samples,  $C_T$  values for all TF genes were normalized to the average  $C_T$  value of the housekeeping genes *UBQ10* (*At4g05320*), *PP2A* (*At1g13320*), *SAND Box Protein* (*At2g28390*) and *GAPDH* (*At1g13440*) (Czechowski *et al.*, 2005), obtaining a  $\Delta C_T$  value for each gene tested in each sample. The TF relative gene expression between aerobic and anaerobic samples was obtained from the equation  $(1 + E)^{-\Delta\Delta C_T}$ , where  $\Delta\Delta C_T$  represents  $\Delta C_{T\text{hypoxia}}$  minus  $\Delta C_{T\text{normoxia}}$ , and  $E$  is the PCR efficiency.

#### Quantification of mature miRNA by stem-loop quantitative PCR (SL-PCR)

The expression of mature miRNAs was assayed by SL-PCR and performed as described previously (Varkonyi-Gasic *et al.*, 2007). The stem-loop reverse transcription primers used have been previously described by Varkonyi-Gasic *et al.* (2007). The reverse transcription reactions were performed on 200 ng total RNA using Superscript III (Invitrogen) and carried out in a pulsed-reverse transcription reaction as described by Varkonyi-Gasic *et al.* (2007). Two cDNA reactions were performed, one containing the miRNA specific primer, and one for the housekeeping control U6, as in Moldovan *et al.* (2009). The reverse transcription product was amplified using a miRNA specific forward primer and a universal reverse primer on a Rotor-Gene 6000 (Invitrogen) in an optical 384-well plate with an ABI PRISM 7900 HT Sequence Detection System (Applied Biosystems), using SYBR Green to monitor dsDNA synthesis. Reactions contained 5  $\mu$ l 2 $\times$  POWER SYBR Green Master Mix reagent (Applied Biosystems), 20 ng cDNA and 250 nmol of each gene-specific primer in a final volume of 5  $\mu$ l. The following standard thermal profile was used for all PCRs: 95°C for 10 min; 40 cycles of

95°C for 15 s and 60°C for 50 s. A final denaturation step (95°C for 15 min, 60°C for 15 min, 95°C for 15 min) was used to examine dissociation curves and test the presence of one single amplicon. Relative RNA levels were calculated using the  $\Delta\Delta C_T$  method as described earlier.

#### Bioinformatic promoter analysis

Differentially expressed genes (DEG) ( $\geq 2$  FC or  $\leq -2$  FC) from the van Dongen *et al.* (2009) dataset (Gene Expression Omnibus accession number: GSE11558; <http://www.ncbi.nlm.nih.gov/geo/>) were selected and split into four groups: induced after 30 min hypoxia (1% and 4% O<sub>2</sub>), repressed after 30 min hypoxia, induced after 2 h hypoxia and repressed after 2 h hypoxia. Sequences of 1000 bp upstream from the transcription start site of the selected genes were downloaded from the Arabidopsis Information Resource (TAIR) database (<http://www.arabidopsis.org/tools/bulk/index.jsp>) using the Bulk Data Retrieval tool. Identification of over-represented hexameric and octameric motifs was carried out using the Motif Analysis (<http://www.arabidopsis.org/tools/bulk/motiffinder/index.jsp>) and by BioProspector Search tool (Liu *et al.*, 2001). Identification of known DNA elements bound by TFs in the promoter dataset was carried out using the web-based analysis tool PlantPAN (Chang *et al.*, 2008) and Pscan (Zambelli *et al.*, 2009), using default parameters.

#### Reporter transactivation assay in protoplasts

Coding sequences (CDSs) of At1g72360, At3g02550, At1g69570, At5g66980 and At4g29190 were amplified from cDNA of hypoxic Arabidopsis roots. *Renilla* luciferase CDS was amplified using pRL-null (Promega) as a template. The resulting amplicons were ligated into the pENTR/D-topo vector (Invitrogen) and subsequently recombined in p2GW7 (Karimi *et al.*, 2002), using the LR clonase enzyme (Invitrogen) to generate the 35S::TF test vector and the 35S::RLuc normalization vector. Arabidopsis plants stably expressing the firefly luciferase CDS under control of the full AtHB1 promoter were used to produce protoplasts according to the protocols described by Yoo *et al.* (2007). Protoplasts were transformed with the test and normalization vectors together using 5  $\mu$ g plasmid DNA each, according to Yoo *et al.* (2007). Nonrecombined p2GW7 was used as a control. Dual luciferase reporter assay was performed using the kit provided by Promega and luminescence was read using a GloMax 20/20 Luminometer (Promega).

#### miRNA target gene prediction

Candidate miR target genes were determined using publicly available prediction algorithms, including miRU (Zhang,


2005), the target search in WMD2 (Ossowski *et al.*, 2008) and the prediction tool in the UEA plant sRNA toolkit (Moxon *et al.*, 2008). The programs were used with their default settings.

## Results

### Hypoxic core genes share conserved motifs in their upstream regions

We analysed the genes that were selected for being differentially expressed in root tissues of *Arabidopsis* seedlings during hypoxia (van Dongen *et al.*, 2009) for the occurrence of overrepresented regulatory motifs. This dataset describes the changes occurring after 30 min, 2 h and 48 h at four different oxygen concentrations (1%, 4%, 8% and 21% O<sub>2</sub>) in darkness. As most changes occurred within the first 2 h of the hypoxic treatments, we selected only those genes that showed at least a twofold change (positive or negative) under hypoxia (both 1% and 4% O<sub>2</sub>) compared with the normoxic control (21% O<sub>2</sub>) after 30 min and 2 h. Most of these genes were also induced at 8% O<sub>2</sub>, although to a smaller extent. The resulting lists are shown in the Supporting Information Tables S1–S4. The overlap between the group of genes induced after 30 min (55 genes) and those upregulated after 2 h (52 genes) was high (45 genes). Conversely, out of 40 genes repressed after 30 min hypoxia only eight were still downregulated after 2 h.

We extracted the first 1000 nucleotides upstream of the annotated translation initiation codon (ATG) for each hypoxia-responsive gene (Tables S1–S4) and scanned them for over-represented DNA elements using various freely available analysis tools. Pscan (Zambelli *et al.*, 2009) and PlantPan (Chang *et al.*, 2008) were used to search for known motifs already reported to be bound by TFs and present in the Jaspar, Transfac and Place databases. Both tools coherently identified the binding site for the *Arabidopsis* Homeobox1 and Homeobox5 (AtHb5) TFs, being enriched in both 30 min and 2 h 'upregulated genes' datasets (Table S5). Pscan reported an enrichment of the homeobox elements AATAATT in the promoters of genes induced and repressed after 30 min, and the AGL3 elements in both upregulated and downregulated genes after 2 h (Table S5). Furthermore, the promoter sequences were analysed for the presence of motifs that were not annotated as regulatory elements before. For this purpose, the 'Motifsearch' tool from the TAIR website and the 'Bioprospector' tool were used. Among these newly identified motifs that were significantly ( $P < 0.001$ ) over-represented in the set of hypoxia induced genes, six elements were present in both upregulated datasets (30 min and 2 h) (Fig. 1). Both prediction tools identified the sequences ATCTAT in the 30 min and 2 h datasets and TTGGAT in the 2 h dataset only (Fig. 1, Tables S6, S7). No common




**Fig. 1** Hexamer motifs overrepresented in hypoxia responsive promoters. Venn diagram depicting the number of DNA elements present in the 1 kb upstream region of induced (UP) and repressed (DOWN) genes after 30 min and 2 h hypoxia (1% and 4% O<sub>2</sub>). A full overview with information about the statistical evaluation of the elements can be found in the Supporting Information, Tables S5–S8.

novel DNA element was identified in the downregulated datasets (Fig. 1 and Tables S8, S9).

### Identification of TFs that are differentially expressed during hypoxia

In order to identify TFs that are differentially expressed by hypoxic conditions, we used a qRT-PCR platform (Czechowski *et al.*, 2002; Barrero *et al.*, 2009) that was successfully used in the past (Scheible *et al.*, 2004; Morcuende *et al.*, 2007; Osuna *et al.*, 2007). The current version of this tool allows the quantification of > 1900 *Arabidopsis* TF gene transcripts. The TF transcript profiling experiment was performed with the same set-up and conditions as described in van Dongen *et al.* (2009) for the microarray analysis to provide maximum comparability between the two datasets. Both the hypoxia-treated plants and the normoxic controls were kept in darkness in order to exclude changes induced by variable factors other than the oxygen concentration such as what could be induced by the shift from light to darkness.

The analysis revealed that mild hypoxic conditions (8% O<sub>2</sub>) induced upregulation of a small number of TF genes after 30 min and 2 h from the hypoxia onset (7 and 11 genes, respectively) (Fig. 2). Lower oxygen concentrations increased the number of upregulated TF after 30 min hypoxia (29 and 24 genes under 4% and 1% O<sub>2</sub>, respectively), with a further increase at 1% O<sub>2</sub> (81 genes). The



**Fig. 2** Number of transcription factors (TFs) being upregulated or downregulated by hypoxia through time. Effect of a stepwise decrease in oxygen concentrations on the number of differentially expressed TFs. Arabidopsis seedlings were grown on vertical sterile plates and incubated at (open triangle) 8%, (open square) 4%, or (open circle) 1% (v : v) oxygen for 30 min and 2 h. The amount of genes is plotted that were at least two-times upregulated (open symbols) or downregulated (shaded symbols).

same trend was observed for downregulated genes, although after 2 h at 1% O<sub>2</sub> the number of downregulated genes was again reduced (Fig. 2). Among the induced TF genes, 21 were already induced after 30 min hypoxia and remained upregulated after 2 h and 10 showed increased transcript levels after 2 h only whereas 12 decreased their expression. Twenty-nine TF genes maintained reduced mRNA levels after 30 min and 2 h (Fig. 2). Eight TFs significantly increased their mRNA level after 30 min and were repressed to the aerobic levels or lower after 2 h hypoxia (Fig. 2).

The full list of differentially expressed TF genes is given in Fig. 3. Twelve TF genes were transiently induced by the hypoxic treatment (1% and 4% O<sub>2</sub>) after 30 min and returned to levels comparable to the aerobic controls or lower after 2 h. By contrast, 17 TF transcripts progressively accumulated after 2 h treatment (Fig. 3a). Among the TFs that have been already characterized, members of the AP2/ERF-type family are the most commonly represented in the set of upregulated TFs (eight members), followed by Zinc-finger (six genes) and bHLH-type TFs (five genes) (Fig. 3b). Transcription factors belonging to the bHLH family are also highly represented among the downregulated genes (10 members), together with members from the bZIP and MYB families (eight members each).

Hypoxia responsive TFs belonging to the auxin response factor (ARF), DNA binding with One Finger (DOF), ERF and Homeobox families potentially bind the 'hypoxic core' promoters


To identify TFs that are supposed to play a role in the early induction of the hypoxic core genes, we searched the plant TF database PLANTTFDB (Guo *et al.*, 2008) for DNA elements specifically bound by the protein families to which

the hypoxia responsive TFs belong. We also integrated the DNA elements characterized by previous studies but not included in PLANTTFDB. Subsequently, we carried out an extensive search for the presence of these DNA elements in the first 1000 base pairs upstream the transcriptional start site of the upregulated genes. Four protein families that included hypoxia-responsive TFs were able to bind DNA elements present in the promoter of at least half of the hypoxic responsive core genes. The ERF binding sites ATCTA (Welsch *et al.*, 2007) and ATATT (Kannangara *et al.*, 2007), and the DOF TF binding element CTTTT(T/A) were present at least once in all the promoters of the selected hypoxia-induced genes (upregulated after 30 min and 2 h) (Table S10). Also, the DNA motif bound by Homeobox proteins, AATAATT, is present in the promoter region of 44 hypoxia-induced genes (Table S10). Finally, the TGTCTC element recognized by the ARF TF family, was found in 24 promoters.

### Hypoxia-responsive TFs can upregulate the expression of hypoxia-induced genes

To verify whether the identified TFs have indeed the potential to regulate the expression of hypoxia responsive genes, we assayed their *trans*-activation capacity on the promoters of two anaerobic genes, Alcohol Dehydrogenase (*ADH*) and Nonsymbiotic Hemoglobin1 (*HBI*). Five hypoxia-induced TFs were selected (*At4g29190*/*LBD41*, *At3g02550*/*HRE1*, *At1g72360*; *At1g69570*; *At5g66980*) from different TF families (Zinc Finger, LBD (Lateral Organ Boundary Domain), ERF, DOF, ARF) according to their predicted interaction with the *ADH* and *HBI* promoters (Fig. 4a). Mesophyll protoplasts were produced from transgenic Arabidopsis leaves expressing the firefly (*Photinus pyralis*) luciferase gene fused to the complete upstream sequence of the *At1g77120* (*ADH*, 905 nucleotides) and *At2g16060* (*HBI*, 369 nucleotides) as flanked by the start codon of either *ADH* or *HBI* and the 3'-untranslated region of the next first upstream gene on the chromosome. Protoplasts were then transiently transformed with vectors carrying the selected TFs under the control of the strong and constitutive 35S promoter of the Cauliflower Mosaic Virus (CaMV). The transformed protoplasts were incubated in the dark before the analysis of the reporter expression to exclude any putative effects of light on the selected hypoxic promoters.

In accordance with our *in silico* prediction, the selected TFs induced the expression of the reporter genes when the promoter contained the relevant binding element. The *HBI* promoter gave significant ( $\geq$  twofold) induction of the reporter gene when *LBD41*, *HRE1*, *At1g69570* and *At5g66980* were overexpressed in the protoplasts. The *ADH* promoter, instead, showed a more modest induction ( $\leq$  1.5-fold) by overexpression of *HRE1*, *At4g29190* and *At5g66980*. Of the three TFs with at least one putative



**Fig. 3** Differentially expressed transcription factors (TFs) under different hypoxic conditions. (a) Heatmap showing the relative RNA level of microRNA primary transcripts (pri-miRNAs) significantly increased or decreased ( $P \leq 0.05$ ) after 30 min or 2 h hypoxia (1%, 4%, or 8% (v/v)  $O_2$ ) with respect to normoxia (21% (v/v)  $O_2$ ). Each treatment was repeated twice and only TFs were selected that showed differential expression in at least two treatments. The fold-change is shown as  $\log_2$ . (b) Pie-chart depicting the distribution of induced and repressed TF genes in protein families.

binding site in the two promoters, only HRE1 and At5g66980 were able to transactivate the expression of both reporter constructs, while At1g69570 only affected the luciferase expression in combination with the *HBI* promoter. In accordance with the prediction, At4g29190 only caused induction in combination with the *ADH* promoter, which contains a bipartite zinc finger binding site AATTAAANNNNNAATTA (Park *et al.* 2007). By contrast, overexpression of *LBD41* only led to transactivation when the reporter gene was fused to the *HBI* promoter, although a putative binding site for the LBD family (CGGCG) was identified only in the *ADH* promoter. Summarizing, it can be concluded that, with one exception, the transactivation assays confirmed the *in silico* prediction of the regulatory TF binding elements.


#### Identification of hypoxia-responsive pri-miRNAs

Moldovan *et al.* (2009) identified various mature miRNAs and their precursors being upregulated or downregulated


after 5 h of near-anoxic (0.1%  $O_2$ ) conditions. In order to test whether this regulation takes place after 30 min or 2 h hypoxia and at various oxygen concentrations, we used the pri-miRNA qRT-PCR platform recently reported by Pant *et al.* (2009). Only one pri-miRNA, miR159c, showed increased expression ( $\log_{FC} \geq 2$ ) in hypoxic conditions after 30 min, whereas 17 pri-miRNAs were downregulated ( $\log_{FC} \leq -2$ ) both at 1% and 4%  $O_2$  concentrations (Fig. 5a). After 2 h the number of upregulated pri-miRNA increased to four, while the downregulated pri-miRNA decreased to eight (Fig. 5a). MiR159c is the most induced sequence after 30 min, although the increased expression does not last long, as after 2 h hypoxia it returned to the aerobic expression levels (Fig. 5b).

#### Specific miRNA biosynthesis takes place at the post-transcriptional level

A previous analysis of mature miRNA in Arabidopsis identified several sequences upregulated after 5 h hypoxia



**Fig. 4** Transactivation assay to test the ability of selected hypoxic transcription factors (TFs) to activate gene expression via interaction with promoter elements from *ADH* or *HB1* *Arabidopsis* protoplast. (a) Schematic representation of the putative binding elements of the selected TF on the *ADH* and *HB1* promoters. The 5' untranslated regions are the tinted part of the bars, the 5' intergenic regions are indicated by the open part of the bars. (b) Regulation of *ADH* and *HB1* expression by transient expression of hypoxia-responsive TFs. Transgenic *Arabidopsis* plants stably transformed with the reporter constructs *ADH::fLuc* or *HB1::fLuc* were used to prepare mesophyll protoplasts. The protoplasts were simultaneously transfected with a P35S:TF effector construct and a P35S:rLUC construct for normalization of the signal. Averaged ( $n \geq 6$ ) normalized fLuc/rLuc activities are plotted relative to the empty vector control. Error bars represent standard deviations. Asterisks indicate significant effects (one-way ANOVA test,  $P < 0.05$ ).



**Fig. 5** Differentially expressed microRNAs (miRNA) under different hypoxic condition. (a) Effect of a stepwise decrease in oxygen concentrations on the number of differentially expressed microRNA primary transcripts (pri-miRNAs). *Arabidopsis* seedlings were grown on vertical sterile plates and incubated at different oxygen concentrations for 30 min and 2 h. Plotted here are the number of genes that were at least two-times up-regulated (open symbols) or down-regulated (shaded symbols) by either (open triangle) 8%, (open square) 4%, or (open circle) 1% (v/v) oxygen at the two different time intervals, as indicated on the horizontal axis. (b) Upregulated and downregulated pri-miRNA after 30 min or 2 h hypoxia. Heatmap showing the relative RNA level of pri-miRNAs significantly increased or decreased ( $P \leq 0.05$ ) after 30 min or 2 h hypoxia (1%, 4%, or 8% (v : v)  $O_2$ ) with respect to normoxia (21% (v : v)  $O_2$ ). The fold-change is shown as  $\log(2)$ .

(Moldovan *et al.*, 2009). By contrast, in our analysis only one pri-miRNA was induced after 30 min hypoxia and three were induced after 2 h. Therefore, we tested the expression of the pri-miRNA reported by Moldovan *et al.* (2009)

and those identified in the present study (pri-miR156g, pri-miR157a, pri-miR158a, pri-miR159a, pri-miR172a, pri-miR172b and pri-miR391) at a more detailed time resolution. we also analysed the level of mature miRNA in parallel. Nevertheless, we could only detect a constant increase of pri-miR391, as already observed in the complete pri-miRNA profiling (Fig. 6). Pri-miR157a and pri-miR172a showed a modest increase after 1 h but decreased again after 2 h and 4 h. Conversely, pri-miR156g remained extremely constant during the hypoxic treatment, while pri-miR158a, pri-miR159a and pri-miR172b were downregulated (Fig. 6).

We tested whether the mature miRNAs, produced from the pri-miRNAs previously analysed, were upregulated or downregulated under hypoxia. Mature miRNA391 showed a modest increase (> twofold) after 4 h, whereas the other mature miRNAs remained constant (miR156, miR157, miR158, miR159, miR172) (Fig. 6). In general, for all the miRNAs examined, changes at the pri-miRNA level did not result in the same magnitude of change in the mature miRNA pool.


#### Putative target transcripts of miRNAs that are differentially expressed under hypoxia do not change their abundance

miRNAs regulate gene expression by endonucleolytic cleavage of target mRNAs and they may also act through inhibition of translation (Bari *et al.*, 2006; Brodersen *et al.*, 2009; Pant *et al.*, 2009). Many of the known targets of miRNA are TF genes. Therefore, we searched for predicted or experimentally determined targets of the miRNA whose pri-miRNA was observed to be upregulated or downregulated under hypoxia (Table S11). The targets of the four upregulated miRNAs identified in this work included MYB TFs (pri-miR159c), protein kinases (pri-miR391), oxidoreductases (miR397) and squamosa promoter binding proteins (SBPs) (miR156e). Analysis of the TF dataset obtained in the present work and the previous whole-transcriptome profiling revealed that none of the targets of the induced pri-miRNA showed a significant downregulation at the transcriptional level within 2 h hypoxia. In a similar fashion, targets of downregulated miRNA did not show induction.

## Discussion

### Hypoxia-specific TFs identified by qRT-PCR

When plants are exposed to hypoxic conditions, they rapidly adapt their transcriptional program to cope with the developing stress conditions (van Dongen *et al.*, 2009). Most of the upregulated mRNAs are also loaded on polysomes to allow their translation (Branco-Price *et al.*, 2008). As TF proteins and small RNAs play a crucial role in the control of the transcriptional machinery, unveiling their



**Fig. 6** Time-course analysis of the effect of hypoxia (1% O<sub>2</sub>) on selected microRNA primary transcripts (pri-miRNAs) and their mature miRNA. Relative RNA levels of miRNAs precursors (pri-miR156a, pri-miR157a, pri-miR158a, pri-miR159a, pri-miR172a, pri-miR172b and pri-miR391, pri-miR169d, pri-miR829, pri-miR395b, pri-miR397b) and mature miRNAs (miR156, miR157, miR158, miR159, miR172, miR391, miR169, miR829, miR395, miR397) in roots of 15-d old seedling of Col-0 plants. Changes of the relative RNA levels were measured during an 8-h time-course by real-time quantitative PCR- calculated according the 2<sup>-ΔΔCT</sup> method and shown as fold-change values (1 = 0 h Hypoxia).

targets and their regulation represent a milestone in the understanding of plant adaptation to oxygen shortage.

The qRT-PCR technique represents a sensitive assay to quantify changes in the abundance of TF and pri-miRNA

transcripts, which are often expressed at levels that are undetectable by microarrays (Czechowski *et al.*, 2002; Scheible *et al.*, 2004; Morcuende *et al.*, 2007; Osuna *et al.*, 2007). Previous microarray analyses identified some TF genes upregulated by low oxygen conditions (Klok *et al.*, 2002; Loreti *et al.*, 2005; van Dongen *et al.*, 2009). Targeted analyses of specific cell populations, such as those performed by Mustroph *et al.* (2009), can circumvent the sensitivity limitation of the microarray technique. The signal for cell specific mRNA is increased by the enrichment derived by the immune-precipitation (Mustroph *et al.*, 2009). However, this study mainly addressed changes in the transcriptome and could not distinguish between changes at the transcript level or in the association of mRNA with polysomes. Furthermore, this technique still suffers of sensitivity limitation for mRNA whose amount is low in all cell types. For all these reasons, the qRT-PCR technique is considered to be optimally suited to address changes in the mRNA level of TFs and pri-miRNAs. Thirteen TF genes were upregulated and 33 TF-genes were downregulated after 30 min of either 1% (v/v) or 4% (v/v) oxygen compared with normoxia (21% v/v oxygen) (Fig. 2). After 2 h treatment, 49 TF transcripts were induced and 23 repressed by hypoxia (Fig. 2). A part of the differentially regulated TF genes found in the current study was not identified by previous microarray analyses. This may be because the corresponding probe set is not present on the chip or no significant change was detected as a result of the relatively low expression levels of TFs. Furthermore, it is possible that the discrepancies with previous studies are caused by the differences in experimental set-up: in particular, the upregulation of genes encoding bHLH and homeobox TFs has not been reported before. Interestingly, a set of 12 genes was transiently induced after 30 min hypoxia. Apart from HEC1 and ABF1 proteins, most of these TFs have not yet been characterized. They might be involved in the initiation of the hypoxic response. The low oxygen responsive genes *ADH* and *HBI* require protein synthesis, most likely of one or more TFs, to be induced under hypoxia (Licausi *et al.*, 2010). The rapid induction of these TF under hypoxia suggests that their regulation depends on release of a repressor or post-transcriptional activation of an activator. Within the group of TF genes induced by hypoxia (Fig. 3a), we also found some that were already reported as hypoxic genes and can be considered as markers of the hypoxic treatment, such as *Hypoxia Responsive ERF1* and 2 (*HRE1* and *HRE2*) (Licausi *et al.*, 2010), LOB (Lateral Organ Boundary) Domain *LBD41* and *At3g10040* (Fig. 3a).

#### Reactive oxygen species (ROS)-related TFs are induced by anoxia but not hypoxia

Not only did we discover new TFs or confirmed already characterized changes in TF expression upon hypoxia with

our experiments, but there were also a few TFs that were not differentially expressed in our approach, although their hypoxia-induced regulation was reported in the literature previously. This is exemplified for heat shock factor *HSFA2*, and the zinc finger proteins *ZAT10* and *ZAT12*, which were not induced in our experiments even at 1% O<sub>2</sub>, although these genes were reported to be induced under by low oxygen treatment by several other studies (Branco-Price *et al.*, 2005, 2008; Loreti *et al.*, 2005). *HSFA2* is involved in the upregulation of several heat shock proteins (HSPs) and other oxidative stress-related proteins under anoxia (Banti *et al.*, 2008) and its over-expression improves anoxia and submergence tolerance in *Arabidopsis* (Banti *et al.*, 2010). Both *ZAT10* and *ZAT12* are also transcription factors that mediate the molecular response to ROS in plants (Davletova *et al.*, 2005; Mittler *et al.*, 2006). As our previous transcriptome analyses (van Dongen *et al.*, 2009; Licausi *et al.*, 2010) also did not show an increase in *HSFA2* and its targets under constant hypoxia, it is likely that the upregulation of ROS-related TFs constitute a specific response to true anoxic conditions in the dark, whereas in our experimental system oxygen concentration never declined below 1%. Indeed, Loreti *et al.* (2005) used anoxic conditions and Branco-Price *et al.* (2005, 2008) flushed an O<sub>2</sub>-free argon atmosphere in a sealed container, reaching complete anoxia after some hours. Apparently, the induction of hypoxic TF genes is (at least partly) regulated by a signaling pathway that is different from the regulation mechanism of anoxic responses.

At mild hypoxic conditions (8% O<sub>2</sub>), the expression of fewer genes was affected compared with more extreme hypoxic conditions (1% v/v oxygen). Nevertheless, some TFs that are strongly induced by strict hypoxia also exhibit upregulation at 8% O<sub>2</sub>. This is likely to be because the intensity of the hypoxic signal is inversely dependent on the oxygen levels within the cell. Considering the induction magnitude under mild and severe hypoxia, two groups of anaerobic genes could be distinguished: those that require more time to be induced, when treated with mild hypoxia (8% O<sub>2</sub>) (such as *At3g10040*, *At3g02550* and *At1g67100*), and those that are not induced by the mild-hypoxic treatment (such as *At2g47520*, *At1g72210*, *At3g16060* and *At1g18400*). It is tempting to speculate that the sensing mechanism and signaling pathways that lead to the induction of these two groups of genes are different.

#### TFs induced by hypoxia bind overrepresented elements in the promoter of anaerobic genes

Individual members of a TF family are suggested to bind different oligonucleotide target sequences, although usually a consensus DNA element can be identified (Jin & Martin, 1999; Toledo-Ortiz *et al.*, 2003; Yang *et al.*, 2009). The promoter sequences of the core set genes were analysed for

overrepresented regulatory sequence motifs. As expected, 5 bp sequences were found more frequently than 6- or 8-bp long ones. We identified several DNA elements that are known to be bound by some ERF, DOF and homeobox TFs (Table S9). The ERF-bound sequence ATATT (Kannangara *et al.*, 2007) is also homologous to part of the TATA-box, which constitutes a common feature of most genes transcribed by the RNA polymerase II (Smale & Kadonaga, 2003). However, the Motifsearch tool identified it as significantly overrepresented with respect to the genes present in the Arabidopsis genome, suggesting that A/T rich regions may have a further role under hypoxia unrelated to the assembly of the RNA polymerase II core complex, as enhancers of transcription.

We confirmed the over-representation of a GC motif (GCCCC) and a GT motif (TGGTTT) in the promoter of hypoxia-induced genes (Fig. 1 and Table S7) as already shown by Liu *et al.* (2005) and Mohanty *et al.* (2005). The GT motif has been proposed to be essential for AtMYB2 binding and hypoxia-induced *ADH1* expression (Hoeren *et al.*, 1998). The *MYB2* gene was not induced by hypoxia either in the microarray (Tables S1, S2) and the qRT-PCR (Table S12) dataset. It is possible that MYB2 is also part of the specific response to anoxia or to an anoxia-derived signal similar to that suggested for as HSF2, ZAT10 and ZAT12.

Another DNA motif, ATCTA, already reported by Welsch *et al.* (2007), was also identified by Motiffinder and Bioproscpector as significantly over-represented in the promoter of upregulated genes with respect to its average occurrence in the Arabidopsis genome. Moreover, this motif has already been reported as conserved in the promoter of anaerobic genes in different plant species (Mohanty *et al.*, 2005). The only TF reported until now to be able to bind this element is the ERF protein AtRAP2.2 (Welsch *et al.*, 2007). Interestingly, AtRAP2.2 was shown to be important for ethylene-mediated tolerance to hypoxia in Arabidopsis seedlings (Hinz *et al.*, 2010). RAP2.2 and its homologs RAP2.12 and RAP2.3 do not respond to hypoxia or anoxia (Loreti *et al.*, 2005; van Dongen *et al.*, 2009; present study). Two other homologs of *RAP2.2*, *HRE1* and *HRE2*, were induced by hypoxia in our dataset (Table S2). These two genes are required to maintain the high expression of fermentative enzymes and indeed, *HRE1* overexpression caused an increase in the upregulation of anaerobic genes under hypoxia (Licausi *et al.*, 2010). Interestingly, when *HRE1* was transiently overexpressed in mesophyll protoplasts, it induced the expression of a reporter gene fused to the promoter of *ADH* or *HB1* genes, which both contain the ATCTA element (Fig. 4).

It is tempting to speculate that the bHLH TFs may also act as mediators in oxygen signaling, as oxygen sensors from animals (HIF; Wenger, 2002), and fungi (SRN1; Goldstein *et al.*, 2006) belong to this family. Our dataset revealed five

bHLH genes being induced by hypoxia in our dataset, although only few genes in the 'hypoxia-induced' cluster (Table S10) contained the bHLH binding element in their promoters. This would rule out the bHLH proteins as regulators of the hypoxic response in Arabidopsis. However, bHLH are known to create homo-oligomer or hetero-oligomer complexes and therefore they may still be involved in the regulation of hypoxic genes, in association with other upregulated TFs. For example, some bHLH proteins have been shown to interact with members of the LBD protein family (Husbands *et al.*, 2007). Interestingly, two members of the LBD family were also upregulated by hypoxia in all the conditions tested in the current experiments (*LBD 40* and *LBD41*). Rubin *et al.* (2009) showed that three highly related *LBD* genes (*LBD37*, *LBD38* and *LBD39*) act as repressors of a larger subset of nitrogen-starvation induced genes. However, in our transactivation assay in protoplasts, the overexpression of *LBD41* caused an upregulation of the reporter gene controlled by the *HB1* promoter. Moreover, the data reported by Wang *et al.* (2003) showed that *HB1* is induced by nitrate and nitrite supplementation, together with *LBD37*, *LBD38* and *LBD39*, suggesting that LBDs may act as positive and negative regulators on different targets, perhaps through the interaction with members of the bHLH family. Analyses of the protein-protein interaction among the upregulated TFs will help in clarifying this aspect.

Pscan and Plantpan identified the binding site of *HOMEBOX1* (*AtHB1*) and *AtHB5* genes as over-represented in the upregulated dataset. These specific TFs are not among the homeobox genes that exhibited upregulation (*AtWAX14*, *At4g33280* and *At4g02550*) by hypoxia in our dataset; however the general homeobox element AATAATT is present in the promoter of 44 anaerobic genes. Homeobox proteins are conserved among all eukaryotes and share basic structural similarity with the helix-loop-helix domains of some bacterial proteins. In plants, these TFs are involved in the regulation of meristem development, especially in stem cell niches (Scofield & Murray, 2006). Severe hypoxic stress, such as the one imposed by submergence, causes death of mature leaves and roots and reduces the size of new organs (Bailey-Serres & Voisenek, 2008). It is possible that the homeobox genes are involved in reprogramming the development of new roots and leaves to adapt to the environmental condition.

In rice, *in silico* analyses identified putative TF and DNA elements involved in the regulation of the molecular response to anaerobiosis also (Narsai *et al.*, 2009). Interestingly, only a little overlap was observed between the motifs over-represented in the promoters of hypoxia responsive genes in Arabidopsis and rice. For example, several elements containing the LBD binding site were over-represented in the promoter of anaerobic genes in rice (Narsai *et al.*, 2009), whereas the hypoxia responsive


promoters do not contain such elements in *Arabidopsis*. This observation suggests that these two plant species possess different subsets of transcription factors and DNA binding elements to modulate their hypoxic response. Nevertheless, several orthologous members of specific families – including the ERF, LBD and trihelix TFs – exist that are coherently responding to hypoxia in several different species and therefore seem to constitute a core element in the response of plants to oxygen deprivation (Loreti *et al.*, 2005; Branco-Price *et al.*, 2005; Lashanti-Kudahettige *et al.*, 2007; Pasentsis *et al.*, 2007; Kreuzwieser *et al.*, 2009; Narsai *et al.*, 2009).

### Hypoxia induces few pri-miRNAs although induction of their mature products was not always observed

Regulation of TF transcript levels may be by other TFs but also by the activity of miRNAs and ta-siRNAs (Shukla *et al.*, 2008). miRNAs have been shown to play important roles in the transduction of the hypoxic signal and the related development in animals (Crosby *et al.*, 2009). We used the qRT-PCR platform designed by Pant *et al.* (2009) to quantify changes in the expression of 177 *Arabidopsis* pri-miRNAs. We could identify only one pri-miRNA up-regulated after 30 min hypoxia and four after 2 h hypoxia

(Fig. 5). Instead, using high-throughput sequencing, Moldovan *et al.* (2009) identified 20 mature miRNA significantly upregulated after 5 h hypoxia (0.1% O<sub>2</sub>). The only similarity between the two datasets is the upregulation of miR391 after 2 h hypoxia. For all the other miRNAs we could not observe an increase in the corresponding pri-miRNAs, as was reported by Moldovan *et al.* (2009) even when we analysed the expression in a detailed time-resolved manner (Fig. 6). Interestingly, the promoter of the gene coding for the pri-miR391 (*AT5G60408*) also contained the ATCTA elements that was identified in our *in silico* analysis.

The expression of pri-miRNAs that belong to the same family did not always reflect the levels of mature miRNAs, as observed for miR391, miR157a and miR172a (Fig. 6). Therefore, it is not surprising that we did not observe changes in the mRNA levels of the putative targets of the miRNAs whose precursor responded to hypoxia. However this cannot rule out the function of these mature miRNAs as translational inhibitors, as shown by Brodersen *et al.* (2008). Alternatively, the hypoxia-responsive miRNAs could also be acting in a tissue-specific manner and therefore the effect on their targets may only be observed in specific cell types. Little overlap was also observed between hypoxia-responsive miRNAs in *Arabidopsis* and submergence-responsive miRNA in maize roots (Zhang *et al.*, 2008;



**Fig. 7** Model of the genetic hypoxia signal transduction pathway in *Arabidopsis* roots. Oxygen deficiency is perceived by means of a yet unknown sensor, which triggers the upregulation of a set of hypoxia-responsive transcription factors (TFs) including members of the ERF (HRE1, HRE2), LBD (LBD40 and LBD41), B3 (At5g66980), homeobox (At3g33280) and DOF (At1g69570) families. Under conditions where the oxygen availability is reduced (hypoxia), the subtle regulation of hypoxia-induced effector genes such as *ADH* or *HB1* or specific miRNAs can be controlled via simultaneous interaction of various combinations of these TFs. Under anoxic conditions, an additional set of TFs is activated, including ZAT10, ZAT12 and HSF2, that regulate the differential expression of genes that are only active when molecular oxygen is absolutely absent from the cell.

Moldovan *et al.*, 2009), suggesting that the miRNA response under hypoxia is species-specific.

### Molecular regulation of the low oxygen response in plants

With this study we provide an overview of expressional changes of TFs and miRNAs as transcriptional regulators under hypoxia. The results are summarized in a functional network depicted in Fig. 7. Different from animal systems, miRNAs do not seem to play a major role in the early phases of the regulation of hypoxic genes in plants. Few pri-miRNAs were significantly upregulated under hypoxia but this increase led to an increase of the corresponding mature miRNAs of miR391 only. Our experiments suggest a differentiation between hypoxia and anoxia-induced signals. Some TFs, including HSFs proteins, previously reported to be upregulated by anaerobiosis ( $[O_2] \cong 0\%$  v/v oxygen), did not exhibit a significant increase in the hypoxic conditions tested in our study ( $[O_2] \geq 1\%$  v/v oxygen), suggesting a differentiation of the signal to regulate the expression of various subsets of hypoxia- and anoxia-induced genes. Apart from the oxygen concentration itself, several related parameters can be involved in the transmission of the low-oxygen signal, such as the energy status (Zabalza *et al.*, 2009), the cellular pH (Dennis *et al.*, 2000), or the level of ROS and RNS (Blokina & Fagerstedt, 2010). As all our hypoxic treatments, as well as the aerobic controls, were conducted in the darkness, we were able to separate the effect of the hypoxic treatment from that of other variables such as the shift from light to darkness shift (Hinz *et al.*, 2010). The reciprocal analysis of DNA motifs over-represented in the promoters of genes upregulated by hypoxia and of those elements bound by hypoxia-induced TF suggested that ERF, B3, homeobox, LBD and DOF proteins play a leading role in the regulation of the core set of hypoxic genes, including *miR391*. Regulation of these TFs appeared to occur in a time-dependent manner, with some TFs having their strongest induction in the very early (30 min) phase after the onset of hypoxia, and others being most strongly regulated after 2 h. Our analysis showed that most hypoxia-regulated genes possess various different TF binding elements in their promoter region. The time-resolved regulation of the expression of these genes during hypoxia is therefore suggested to result from the integration of TF-induced gene activation of various TFs binding to different promoter elements simultaneously.

### Acknowledgements

We thank Susanne Freund and Dirk Hinch for maintenance of the transcription factor RT-qPCR platform. This work was supported by the Deutsche Forschungsgemeinschaft (grants nos. Do1298/2-1, Ge878/5-1 and Ge878/7-1).

### References

- Armstrong W. 1979. Aeration in higher plants. *Advances in Botanical Research* 7: 225–332.
- Aukerman MJ, Sakai H. 2003. Regulation of flowering time and floral organ identity by a microRNA and its APETALA2-like target genes. *The Plant Cell* 15: 2730–2741.
- Bailey-Serres J, Voisenek LACJ. 2008. Flooding stress: acclimations and genetic diversity. *Annual Reviews of Plant Biology* 59: 313–339.
- Banti V, Loreti E, Novi G, Santaniello A, Alpi A, Perata P. 2008. Heat acclimation and cross-tolerance against anoxia in Arabidopsis. *Plant, Cell & Environment* 31: 1029–1037.
- Banti V, Mafessoni F, Loreti E, Alpi A, Pierdomenico Perata P. 2010. The heat-inducible transcription factor HSFA2 enhances anoxia tolerance in Arabidopsis. *Plant Physiology* 152: 1471–1483.
- Bari R, Pant DB, Stitt M, Scheible WR. 2006. PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants. *Plant Physiology* 141: 988–999.
- Barrera JM, Millar AA, Griffiths J, Czechowski T, Scheible WR, Udvardi M, Reid JB, Ross JJ, Jacobsen JV, Gubler F. 2009. Gene expression profiling identifies two regulatory genes controlling dormancy and ABA sensitivity in Arabidopsis seeds. *Plant Journal* 61: 611–622.
- Blokina O, Fagerstedt KV. 2010. Oxidative metabolism, ROS and NO under oxygen deprivation. *Plant Physiology and Biochemistry* 48: 359–373.
- Bond DM, Wilson IW, Dennis ES, Pogson BJ, Jean Finnegan E. 2009. Vernalization insensitive 3 (VIN3) is required for the response of *Arabidopsis thaliana* seedlings exposed to low oxygen conditions. *Plant Journal* 59: 576–587.
- Borisjuk L, Macherel D, Benamar A, Wobus U, Rolletshek H. 2007. Low oxygen sensing and balancing in plant seeds: a role for nitric oxide. *New Phytologist* 176: 813–823.
- Branco-Price C, Kaiser KA, Jang CJH, Larive CK, Bailey-Serres J. 2008. Selective mRNA translation coordinates energetic and metabolic adjustments to cellular oxygen deprivation and reoxygenation in *Arabidopsis thaliana*. *Plant Journal* 56: 743–755.
- Branco-Price C, Kawaguchi R, Ferreira RB, Bailey-Serres J. 2005. Genome-wide analysis of transcript abundance and translation in Arabidopsis seedlings subjected to oxygen deprivation. *Annals of Botany* 96: 647–660.
- Branscheid A, Sieh D, Pant BD, May P, Devers E, Elkrog A, Schauer L, Scheible WR, Kranjinski F. 2010. Expression pattern suggest a role of miR399 in the regulation of the cellular response to local Pi-increase during arbuscular mycorrhizal symbiosis. *Molecular Plant and Microbe Interactions* 23: 915–926.
- Broderson P, Sakvarelidze-Achard L, Bruun-Rasmussen M, Dunoyer P, Yamamoto YY, Sieburth L, Voinnet O. 2008. Widespread translational inhibition by plant miRNAs and siRNAs. *Science* 320: 1185–1190.
- Caldana C, Scheible WR, Mueller-Roeber B, Ruzicic S. 2007. A quantitative RT-PCR platform for high-throughput expression profiling of 2500 rice transcription factors. *Plant Methods* 3: 3–7.
- Chang WC, Lee TY, Huang HD, Huang HY, Pan RL. 2008. PlantPAN: Plant promoter analysis navigator, for identifying combinatorial cis-regulatory elements with distance constraint in plant gene groups. *BMC Genomics* 9: 561. doi:10.1186/1471-2164-9-561.
- Chiou TJ, Aung K, Lin SI, Wu CC, Chiang SF, Su CL. 2006. Regulation of phosphate homeostasis by microRNA in Arabidopsis. *The Plant Cell* 18: 414–421.
- Christianson JA, Wilson IW, Llewellyn DJ, Dennis ES. 2009. The low-oxygen-induced NAC domain transcription factor ANAC102 affects viability of Arabidopsis seeds following low-oxygen treatment. *Plant Physiology* 149: 1724–1738.
- Crosby ME, Kulshreshtha R, Ivan M, Glazer PM. 2009. MicroRNA regulation of DNA repair gene expression in hypoxic stress. *Cancer Research* 69: 1221–1229.

- Czechowski T, Bari RP, Stitt M, Scheible W-R, Udvardi MK. 2002. Real-time RT-PCR profiling of over 1400 Arabidopsis transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific genes. *Plant Journal* 38: 366–379.
- Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR. 2005. Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiology* 139: 5–17.
- Davletova S, Rizhsky L, Liang H, Shengqiang Z, Oliver DJ, Coutu J, Shulaev V, Schlauch K, Mittler R. 2005. Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of Arabidopsis. *The Plant Cell* 17: 268–281.
- Dennis ES, Dolferus R, Ellis M, Rahman M, Wu Y, Hoeren FU, Grover A, Ismond KP, Good AG, Peacock WJ. 2000. Molecular strategies for improving waterlogging tolerance in plants. *Journal of Experimental Botany* 51: 89–97.
- van Dongen JT, Froehlich A, Ramírez-Aguilar SJ, Schauer N, Fernie AR, Erban A, Kopka J, Clark J, Langer A, Geigenberger P. 2009. Transcript and metabolite profiling of the adaptive response to mild decreases in oxygen concentration in the roots of Arabidopsis plants. *Annals of Botany* 103: 269–280.
- van Dongen JT, Roeb GW, Dautzenberg M, Froelich A, Vigeolas H, Minchin PEH, Geigenberger P. 2004. Phloem import and storage metabolism are highly coordinated by the low oxygen concentrations within developing wheat seeds. *Plant Physiology* 135: 1809–1821.
- van Dongen JT, Schurr U, Pfister M, Geigenberger P. 2003. Phloem metabolism and function have to cope with low internal oxygen. *Plant Physiology* 131: 1529–1543.
- Emery JF, Floyd SK, Alvarez J, Eshed Y, Hawker NP, Izhaki A, Baum SF, Bowman JL. 2003. Radial patterning of Arabidopsis shoots by class III HD-ZIP and KANADI genes. *Current Biology* 13: 1768–1774.
- Forristal CE, Wright KL, Hanley NA, Oreffo RO, Houghton FD. 2010. Hypoxia inducible factors regulate pluripotency and proliferation in human embryonic stem cells cultured at reduced oxygen tensions. *Reproduction* 139: 85–97.
- Geigenberger P. 2003. Response of plant metabolism to too little oxygen. *Current Opinion in Plant Biology* 6: 247–256.
- Goldstein JL, Debose-Boyd RA, Brown MS. 2006. Protein sensors for membrane sterols. *Cell* 124: 35–46.
- Guo AY, Xin Chen X, Gao G, Zhang H, Zhu QH, Liu XC, Zhong YF, Gu X, He K, Luo J. 2008. PlantTFDB: a comprehensive plant transcription factor database. *Nucleic Acids Research* 36: 966–969.
- Hinz M, Wilson IW, Yang J, Buerstenbinder K, Llewellyn D, Dennis ES, Sauter M, Dolferus R. 2010. Arabidopsis RAP2.2: an ethylene response transcription factor that is important for hypoxia survival. *Plant Physiology* 153: 757–772.
- Hoeren FU, Dolferus R, Wu Y, Peacock WJ, Dennis ES. 1998. Evidence for a role for AtMYB2 in the induction of the Arabidopsis Alcohol Dehydrogenase gene (ADH1). *Low Oxygen Genetics* 149: 479–490.
- Huang SQ, Xiang AL, Che LL, Chen S, Li H, Song JB, Yang ZM. 2010. A set of miRNAs from *Brassica napus* in response to sulphate deficiency and cadmium stress. *Plant Biotechnology Journal*. doi: 10.1111/j.1467-7652.2010.00517.x
- Husbands A, Elizabeth M, Bell EM, Shuai B, Smith HM, Springer PS. 2007. Lateral organ boundaries defines a new family of DNA-binding transcription factors and can interact with specific bHLH proteins. *Nucleic Acids Research* 35: 6663–6671.
- Ismond KP, Dolferus R, De Pauw M, Dennis ES, Good AG. 2003. Enhanced low oxygen survival in Arabidopsis through increased metabolic flux in the fermentative pathway. *Plant Physiology* 132: 1292–1302.
- Jin H, Martin C. 1999. Multifunctionality and diversity within the plant MYB-gene family. *Plant Molecular Biology* 41: 577–585.
- Kannangara R, Branigan C, Liu Y, Penfielda T, Raob V, Mouillec G, Höftc H, Pauly M, Riechmann JL, Brouna P. 2007. The transcription factor WIN1/SHN1 regulates cutin biosynthesis in *Arabidopsis thaliana*. *The Plant Cell* 19: 1278–1294.
- Karimi M, Inze D, Depicker A. 2002. Gateway vectors for *Agrobacterium*-mediated plant transformation. *Trends on Plant Science* 7: 193–195.
- Klok EJ, Wilson IW, Wilson D, Chapman SC, Ewing RM, Somerville SC, Peacock WJ, Dolferus R, Dennis ES. 2002. Expression profile analysis of the low-oxygen response in Arabidopsis root cultures. *The Plant Cell* 14: 2481–2494.
- Kreuzwieser J, Hauberg J, Howell KA, Adam Carroll A, Rennenberg H, Millar HA, Whelan J. 2009. Differential response of gray poplar leaves and roots underpins stress adaptation during hypoxia. *Plant Physiology* 149: 461–473.
- Lanet E, Delannoy E, Sormani R, Floris M, Brodersen P, Crété P, Voinnet O, Robaglia C. 2009. Biochemical evidence for translational repression by Arabidopsis microRNAs. *The Plant Cell* 21: 1762–1768.
- Lasanthi-Kudahettige R, Magneschi L, Loreti E, Gonzali S, Licausi F, Novi G, Beretta O, Vitulli F, Alpi A, Perata P. 2007. Transcript profiling of the anoxic rice coleoptile. *Plant Physiology* 144: 218–231.
- Licausi F, Perata P. 2009. Low oxygen signaling and tolerance in plants. *Advances in Botanical Research* 50: 139–198.
- Licausi F, van Dongen JT, Giuntoli B, Novi G, Santaniello A, Geigenberger P, Perata P. 2010. HRE1 and HRE2, two hypoxia-inducible ethylene response factors, affect anaerobic responses in *Arabidopsis thaliana*. *Plant Journal* 62: 302–315.
- Liu F, VanToai T, Moy LP, Bock G, Linford LD, Quackenbush J. 2005. Global transcription profiling reveals comprehensive insights into hypoxic response in Arabidopsis. *Plant Physiology* 137: 1115–1129.
- Liu X, Brutlag DL, Liu JS. 2001. BioProspector: discovering conserved DNA motifs in upstream regulatory regions of co-expressed genes. *Pacific Symposium Biocomputing* 2001: 127–138.
- Loreti E, Poggi A, Novi G, Alpi A, Perata P. 2005. A Genome-wide analysis of the effects of sucrose on gene expression in Arabidopsis seedlings under anoxia. *Plant Physiology* 137: 1130–1138.
- Mittler R, Kim Y, Song L, Coutu J, Coutu A, Ciftci-Yilmaz S, Lee H, Stevenson B, Zhu JK. 2006. Gain- and loss-of-function mutations in Zat10 enhance the tolerance of plants to abiotic stress. *FEBS Letters* 580: 6537–6542.
- Mohanty B, Krishnan SPT, Swarup S, Bajic VB. 2005. Detection and preliminary analysis of motifs in promoters of anaerobically induced genes of different plant species. *Annals of Botany* 96: 669–681.
- Moldovan D, Spriggs A, Yang J, Pogson BJ, Dennis ES, Wilson IW. 2009. Hypoxia-responsive microRNAs and *trans*-acting small interfering RNAs in Arabidopsis. *Journal of Experimental Botany* 61: 165–177.
- Morcuende R, Bari R, Gibon Y, Zheng W, Pant BD, Bläsing O, Usadel B, Czechowski T, Udvardi MK, Stitt M *et al.* 2007. Genome-wide reprogramming of metabolism and regulatory networks of Arabidopsis in response to phosphorus. *Plant, Cell & Environment* 30: 85–112.
- Moxon S, Jing R, Szitty G, Schwach F, Rusholme P, Pilcher RL, Moulton V, Dalmay T. 2008. Deep sequencing of tomato short RNAs identifies microRNAs targeting genes involved in fruit ripening. *Genome Research* 18: 1602–1609.
- Mustroph A, Lee SC, Oosumi T, Zanetti MA, Yang H, Ma K, Yaghoobi-Masihi A, Fukao T, Bailey-Serres J. 2010. Cross-Kingdom comparison of transcriptomic adjustments to low-oxygen stress highlights conserved and plant-specific responses. *Plant Physiology* 152: 1484–1500.
- Mustroph A, Zanetti ME, Jang CJ, Holtan HE, Repetti PP, Galbraith DW, Girke T, Bailey-Serres J. 2009. Profiling transcriptomes of discrete cell population s resolves altered cellular priorities during hypoxia in Arabidopsis. *Proceedings of the National Academy of Sciences, USA* 106: 18843–18848.
- Narsai R, Howell KA, Carroll A, Ivanova A, Millar AH, Whealan J. 2009. Defining core metabolic and transcriptomic responses to oxygen

- availability in rice embryos and young seedlings. *Plant Physiology* 151: 306–322.
- Ossowski S, Schwab R, Weigel D. 2008. Gene silencing in plants using artificial microRNAs and other small RNAs. *Plant Journal* 53: 674–690.
- Osuna D, Usadel B, Morcuende R, Gibon Y, Bläsing OE, Höhne M, Günter M, Kamlage B, Trethewey R, Scheible WR *et al.* 2007. Temporal responses of transcripts, enzyme activities and metabolites after adding sucrose to carbon-deprived Arabidopsis seedlings. *Plant Journal* 49: 463–491.
- Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D. 2003. Control of leaf morphogenesis by microRNAs. *Nature* 425: 257–263.
- Pant BD, Buhtz A, Kehr J, Scheible WR. 2008. MicroRNA399 is a long-distance signal for the regulation of plant phosphate homeostasis. *Plant Journal* 53: 731–738.
- Pant BD, Musialak-Lange M, Nuc P, May P, Buhtz A, Kehr J, Walther D, Scheible WR. 2009. Identification of nutrient-responsive Arabidopsis and rapeseed microRNAs by comprehensive real-time polymerase chain reaction profiling and small RNA sequencing. *Plant Physiology* 150: 1541–1555.
- Park HC, Kim ML, Lee SM, Bahk JD, Yun D-J, Lim CO, Hong JC, Lee SY, Cho MJ, Chung WS. 2007. Pathogen-induced binding of the soybean zinc finger homeodomain proteins GmZF-HD1 and GmZF-HD2 to two repeats of ATTA homeodomain binding site in the calmodulin isoform 4 (GmCam4) promoter. *Nucleic Acids Research* 35: 3612–3623.
- Pasentis K, Falara V, Pateraki I, Gerasopoulos D, Kanellis AK. 2007. Identification and expression profiling of low oxygen regulated genes from *Citrus flavedo* tissues using RT-PCR differential display. *Journal of Experimental Botany* 58: 2203–2216.
- Rocha M, Sodek L, Licausi F, Hameed MW, Dornelas MC, van Dongen JT. 2010. Analysis of alanine aminotransferase in various organs of soybean (*Glycine max*) and in dependence of different nitrogen fertilisers during hypoxic stress. *Amino Acids*. doi: 10.1007/s00726-010-0596-1.
- Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP. 2002. Prediction of plant microRNA targets. *Cell* 110: 513–520.
- Rubin G, Tohge T, Matsuda F, Saito K, Scheible WR. 2009. Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in Arabidopsis. *The Plant Cell* 21: 3567–3584.
- Scheible WR, Morcuende R, Czechowski T, Fritz C, Osuna D, Palacios-Rojas N, Schindelasch D, Thimm O, Udvardi MK, Stitt M. 2004. Genome-wide reprogramming of primary and secondary metabolism, protein synthesis, cellular growth processes, and the regulatory infrastructure of Arabidopsis in response to nitrogen. *Plant Physiology* 136: 2483–2499.
- Scofield S, Murray JAH. 2006. KNOX gene function in plant stem cell niches. *Plant Molecular Biology* 60: 929–946.
- Shukla LI, Chinnusamy V, Sunkar R. 2008. The role of microRNAs and other endogenous small RNAs in plant stress responses. *Biochimica et Biophysica Acta* 1779: 743–748.
- Smale ST, Kadonaga JT. 2003. The RNA polymerase II core promoter. *Annual Review of Biochemistry* 72: 449–479.
- Sunkar R, Zhu JK. 2004. Novel and stress-regulated microRNAs and other small RNAs from Arabidopsis. *The Plant Cell* 16: 2001–2019.
- Toledo-Ortiz G, Huq E, Quail PH. 2003. The Arabidopsis Basic/Helix–Loop–Helix transcription factor family. *The Plant Cell* 15: 1749–1770.
- Varkonyi-Gasic E, Wu R, Wood M, Walton EF, Hellens RP. 2007. A highly sensitive RT-PCR method for detection and quantification of microRNAs. *Plant Methods* 3: 12.
- Vigeolas H, van Dongen JT, Waldek P, Huhn D, Geigenberger P. 2003. Lipid storage metabolism is limited by the prevailing low oxygen concentrations within developing seeds of oilseed rape. *Plant Physiology* 133: 2048–2060.
- Wang R, Okamoto M, Xing X, Crawford NM. 2003. Microarray analysis of the nitrate response in Arabidopsis roots and shoots reveals over 1,000 rapidly responding genes and new linkages to glucose, trehalose-6-phosphate, iron, and sulfate metabolism. *Plant Physiology* 132: 556–567.
- Welsch R, Maass D, Voegel T, Della Penna D, Beyer P. 2007. Transcription factor RAP22 and its interacting partner SINAT2: stable elements in the carotenogenesis of Arabidopsis leaves. *Plant Physiology* 145: 1073–1085.
- Wenger RH. 2002. Cellular adaptation to hypoxia: O<sub>2</sub>-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O<sub>2</sub>-regulated gene expression. *FASEB Journal* 16: 1151–1162.
- Yang S, Wang S, Liu X, Yu Y, Yue L, Wang X, Hao D. 2009. Four divergent Arabidopsis ethylene-responsive element-binding factor domains bind to a target DNA motif with a universal CG step core recognition and different flanking bases preference. *FEBS Journal* 276: 7177–7186.
- Yoo SD, Cho YH, Sheen J. 2007. Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nature Protocols* 2: 1565–1572.
- Zabalza A, van Dongen JT, Froelich A, Oliver S, Faix B, Kapuganti JG, Schmalzlin E, Igal M, Orcaray L, Royuela M *et al.* 2009. Regulation of respiration and fermentation to control the plant internal oxygen concentration. *Plant Physiology* 149: 1087–1098.
- Zambelli F, Pesole G, Pavesi G. 2009. Pscan: finding over-represented transcription factor binding site motifs in sequences from co-regulated or co-expressed genes. *Nucleic Acids Research* 37: W247–W252.
- Zhang Y. 2005. miRU: an automated plant miRNA target prediction server. *Nucleic Acids Research* 33: W701–W714.
- Zhang Z, Wei L, Zou X, Tao Y, Liu Z, Zheng Y. 2008. Submergence-responsive microRNAs are potentially involved in the regulation of morphological and metabolic adaptations in maize root cells. *Annals of Botany* 102: 509–519.

## Supporting Information

Additional supporting information may be found in the online version of this article.

**Table S1** List of genes upregulated ( $\geq$  twofold) after 30 min hypoxia (1% and 4% O<sub>2</sub>)

**Table S2** List of genes upregulated after 2 h hypoxia (1% and 4% O<sub>2</sub>)

**Table S3** List of genes downregulated after 30 min hypoxia (1% and 4% O<sub>2</sub>)

**Table S4** List of genes downregulated after 2 h hypoxia (1% and 4% O<sub>2</sub>)

**Table S5** 6-Mer over-represented motif in the promoter of genes in Table S1

**Table S6** 6-Mer over-represented motif in the promoter of genes in Table S2

**Table S7** 6-Mer over-represented motif in the promoter of genes in Table S3

**Table S8** 6-Mer over-represented motif in the promoter of genes in Table S4

**Table S9** Known DNA elements bound by transcription factors (TFs) in the promoter dataset (Tables S1–S4)

**Table S10** DNA binding element of the differentially expressed transcription factors (TFs) under hypoxia and its occurrence in the promoter regions of low oxygen responsive genes (Tables S1, S2)

**Table S11** Expression of the putative targets of differentially expressed microRNA primary transcripts (pri-miRNAs) in the TF dataset and previous microarray analyses

**Table S12** Relative mRNA level of transcription factors (TFs) genes and microRNA primary transcripts (pri-miRNAs) in Arabidopsis roots under different hypoxic conditions

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



### About *New Phytologist*

- *New Phytologist* is owned by a non-profit-making **charitable trust** dedicated to the promotion of plant science, facilitating projects from symposia to open access for our Tansley reviews. Complete information is available at [www.newphytologist.org](http://www.newphytologist.org).
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as-ready' via *Early View* – our average submission to decision time is just 29 days. Online-only colour is **free**, and essential print colour costs will be met if necessary. We also provide 25 offprints as well as a PDF for each article.
- For online summaries and ToC alerts, go to the website and click on 'Journal online'. You can take out a **personal subscription** to the journal for a fraction of the institutional price. Rates start at £149 in Europe/\$276 in the USA & Canada for the online edition (click on 'Subscribe' at the website).
- If you have any questions, do get in touch with Central Office ([newphytol@lancaster.ac.uk](mailto:newphytol@lancaster.ac.uk); tel +44 1524 594691) or, for a local contact in North America, the US Office ([newphytol@ornl.gov](mailto:newphytol@ornl.gov); tel +1 865 576 5261).