

## Rapid report

# Accumulation of anthocyanins in tomato skin extends shelf life

Author for correspondence:

Cathie Martin

Tel: +44 1603 450275

Email: [cathie.martin@jic.ac.uk](mailto:cathie.martin@jic.ac.uk)

Received: 1 July 2013

Accepted: 21 August 2013

Laura Bassolino<sup>1\*</sup>, Yang Zhang<sup>2\*</sup>, Henk-jan Schoonbeek<sup>2</sup>, Claudia Kiferle<sup>1</sup>, Pierdomenico Perata<sup>1</sup> and Cathie Martin<sup>2</sup>

<sup>1</sup>Plant Lab, Scuola Superiore Sant'Anna, Piazza Martiri della Libertà 33, 56127, Pisa, Italy; <sup>2</sup>John Innes Centre, Norwich Research Park,

NR4 7UH, Norwich, UK

*New Phytologist* (2013)

doi: 10.1111/nph.12524

**Key words:** *Aft/Aft atv/atv*, anthocyanins, *Botrytis cinerea*, shelf life, *Solanum lycopersicum*, tomato.

## Summary

- Shelf life is one of the most important traits for the tomato (*Solanum lycopersicum*) industry. Two key factors, post-harvest over-ripening and susceptibility to post-harvest pathogen infection, determine tomato shelf life.

- Anthocyanins accumulate in the skin of *Aft/Aft atv/atv* tomatoes, the result of introgressing alleles affecting anthocyanin biosynthesis in fruit from two wild relatives of tomato, which results in extended fruit shelf life. Compared with ordinary, anthocyanin-less tomatoes, the fruits of *Aft/Aft atv/atv* keep longer during storage and are less susceptible to *Botrytis cinerea*, a major tomato pathogen, post-harvest.

- Using genetically modified tomatoes over-producing anthocyanins, we confirmed that skin-specific accumulation of anthocyanins in tomato is sufficient to reduce the susceptibility of fruit to *Botrytis cinerea*.

- Our data indicate that accumulation of anthocyanins in tomato fruit, achieved either by traditional breeding or genetic engineering can be an effective way to extend tomato shelf life.

## Introduction

Shelf-life is one of the most important agronomic traits for tomato (*Solanum lycopersicum*) and is determined by two components, fruit softening during over-ripening and susceptibility to opportunistic pathogens. *Botrytis cinerea*, better known as gray mold, is the second most important fungal pathogen of plants, economically (Dean *et al.*, 2012). *Botrytis cinerea* can infect vegetables (cabbage, lettuce and broccoli) and fruit crops (grape, red fruit and tomato), as well as a large number of shrubs, trees, flowers, and weeds (Williamson *et al.*, 2007). Several different strategies have been employed to extend tomato shelf life. One major target has been cell-wall modifying enzymes, and different strategies have been developed to decrease their activity (Brummell & Harpster, 2001). Other studies have been directed at increasing the production of antioxidants such as polyamines, because their accumulation is associated with extended shelf life (Valero *et al.*, 2002). The ethylene burst is the key event signaling the onset of ripening in climacteric fruits such as tomato. Manipulation of ethylene biosynthesis and signaling has resulted in varieties with delayed ripening (Vicente *et al.*, 2007).

\*These authors contributed equally to this work.

However, all attempts have resulted in only modest delays to the fruit softening processes and are often accompanied by reduced flavor, texture and aroma of tomato fruit (Vicente *et al.*, 2007). Anthocyanins are a group of natural pigments, widely distributed in most vascular plants (Grotewold, 2006). They are stress responsive compounds, used for pollinator and dispersor attraction, but they are also important phytonutrients in a healthy diet, having anti-tumor, pro-apoptotic, anti-oxidative, anti-inflammatory and anti-neurodegenerative properties (Buer *et al.*, 2010; De Pascual-Teresa *et al.*, 2010; Spencer, 2010). Due to their dietary health benefits, anthocyanins are often targets for engineering and plant-breeding programs. Crops having sub-optimal concentration of anthocyanins, like tomato, have been genetically modified to increase their content (Butelli *et al.*, 2008; Gonzali *et al.*, 2009). Several mutants of tomato, altered in their ability to synthesize anthocyanins have been described (Al-sane *et al.*, 2011). The dominant gene *Aft* (*Anthocyanin fruit*) derived from the interspecific cross of *Solanum lycopersicum* (tomato) to *S. chilense*, shows anthocyanin production in the skin of fruit (Jones *et al.*, 2003). *Aft* triggers anthocyanin production and accumulation in fruits upon stimulation by high light (Mes *et al.*, 2008). The *Aft* gene has been suggested to encode a MYB-related transcription factor (Sapir

*et al.*, 2008). A recessive gene, *atv* (*atrovioleacea*), was introgressed into domesticated tomato plants following a cross between *S. lycopersicum* and *S. cheesmaniae* (L. Riley) Fosberg, and influences anthocyanin pigmentation in the entire tomato plant, particularly in stems and leaves (Mes *et al.*, 2008). Tomato plants homozygous for both *Aft* and *atv* alleles show intensely purple-pigmented fruits (Mes *et al.*, 2008). Anthocyanin synthesis in *Aft/Aft atv/atv* is stimulated significantly by high light and is limited to the epidermis and the pericarp of the fruit, which may have both purple and red regions, depending on exposure of the fruit to light (Supporting Information, Fig. S1). Recently, we reported that purple tomatoes, producing anthocyanin throughout the fruit as a result of the ectopic expression of Delila and Rosea1 transcription factors from *Antirrhinum majus*, have double the shelf life of controls (Zhang *et al.*, 2013). In this study, we show that the accumulation of anthocyanins in *Aft/Aft atv/atv* tomatoes, which is predominantly in the skin, is also associated with extended shelf life. Our finding has important agronomic and commercial implications, since *Aft/Aft atv/atv* tomatoes are naturally enriched in anthocyanins and have extended shelf life.

## Materials and Methods

### Storage tests

Near isogenic lines for either *Aft/Aft* or *atv/atv* mutations are not available, so *Solanum lycopersicum* L. cv Ailsa Craig was chosen as a control tomato line for all the analyses. This choice was made because, unlike the *Aft* and *atv* mutant lines, Ailsa Craig does not produce anthocyanins in the skin of fruit although it shows the same vegetative and fruit characteristics (morphology of the plant and its fruit, size of mature tomatoes, and fruit ripening time) compared with *Aft/Aft*, *atv/atv* and *Aft/Aft atv/atv* fruit (Povero *et al.*, 2011).

Wild-type (WT) (cv Ailsa Craig) and *Aft/Aft atv/atv* fruit were tagged at breaker (when the color of WT fruit and the low-anthocyanin regions of *Aft/Aft atv/atv* fruit begin to turn yellow). To induce high anthocyanin production in *Aft/Aft atv/atv* fruit, tomatoes were grown with supplemented light. *Aft/Aft atv/atv* fruit grown under high light have strong, uniform anthocyanin accumulation in the skin (Supporting Information, Fig. S1) (Mes *et al.*, 2008; Povero *et al.*, 2011). Fruit were harvested at 7 d post-breaker (d0 = 7 dpb). All fruits were sterilized in 10% bleach for 10 min, followed by rinsing in sterilized water and air-drying. Each fruit was placed in a plastic jar and kept at 17°C or at room temperature (RT) under light. Every week, the fresh weight of each fruit was measured and visual softening and collapse of the fruit were assessed (Nambeesan *et al.*, 2010). After measurement, fruit were transferred to a new jar.

### TEAC assay and anthocyanin quantification

TEAC (Trolox equivalent antioxidant capacity) analysis of *Aft/Aft atv/atv* tomatoes was performed at breaker as described by (Pellegrini *et al.*, 2003). Results were expressed as TEAC in mmol of Trolox per kg of fresh weight. Anthocyanin extraction from the

skin of *PRD* tomatoes was performed as described by Butelli *et al.* (2008).

### Measurements of cuticle thickness

Cuticle thickness measurements were modified from the methods described by Yeats *et al.* (2012). WT Ailsa Craig, *Aft/Aft atv/atv* red regions and *Aft/Aft atv/atv* purple regions were sliced into 10–30 µm thick sections, stained with Sudan red (Fluka) (Buda *et al.*, 2009) and thickness was determined using a Leica DM6000 microscope, taking the average of 8 to 10 measurements. The average and standard error of the mean of three biological replicates are reported.

### *Botrytis cinerea* infection

*Botrytis cinerea* (B05.10) was grown and collected as described by Stefanato *et al.* (2009). WT (Ailsa Craig) and *Aft/Aft atv/atv* tomatoes were harvested 14 d after breaker and surface sterilized. Intact WT and *Aft/Aft atv/atv* fruits were sprayed thoroughly with spores ( $2.5 \times 10^5$  spores ml<sup>-1</sup>) three times in the flow cabinet and kept at 20°C, in high humidity. Infection symptoms were observed at 4 d post-inoculation (dpi). For wound inoculation, the fungal culture was diluted with medium to  $5 \times 10^4$  spores ml<sup>-1</sup> (for fruit in the MicroTom genetic background) or  $2.5 \times 10^5$  spores ml<sup>-1</sup> (for WT Ailsa Craig and *Aft/Aft atv/atv* fruits) and incubated at RT for 1.5 h before inoculation. The spore inoculum (5 µl) was added to each wound of both red and purple regions of *Aft/Aft atv/atv* fruits grown under natural light. Lesion diameter was measured 72 h after inoculation. To quantify *B. cinerea* growth using quantitative polymerase chain reaction (qPCR), 1 cm samples of infected fruit tissues were harvested 3 d after inoculation. Seeds were removed and samples were freeze dried. Total DNA was isolated and qPCR was performed as described previously (Zhang *et al.*, 2013).

### Plasmid construction and tomato transformation

The light-responsive, PLI promoter which is active predominantly in fruit peel was kindly provided by Dr Diego Orzaez (Estornell *et al.*, 2009). Using Gateway recombination, the PLI promoter was introduced into pDONR 207 to create pENTR-PLI. The PLI promoter was then inserted into the binary vector pJAM1890 (GATEWAY:Ros1/35S:Del) (Martin *et al.*, 2012) to create pPLI:Ros1/35S:Del (pPRD). pPRD was transferred to *Agrobacterium tumefaciens* strain AGL1 by triparental mating. Tomato variety MicroTom was transformed by dipping cotyledons (Fillatti *et al.*, 1987). More than 40 *PRD* T0 independent transgenic lines were produced. Among these, 12 stable T1 lines accumulating different amounts of anthocyanins were selected for further analysis.

### Staining of seed for proanthocyanidins

Tomato seed were stained for proanthocyanidins using 4-dimethylaminocinnamaldehyde (DMACA) as described previously (Abeynayake *et al.*, 2011).

Statistics

Paired or unpaired, two-tailed Student's *t*-tests were used to compare group differences. *P* values < 0.05 were considered significant.

Results

*Aft/Aft atv/atv* tomato can be stored longer


To test whether softening is delayed in *Aft/Aft atv/atv* tomatoes, we performed storage tests under different conditions. WT (Ailsa Craig) and *Aft/Aft atv/atv* tomatoes (grown with supplemental light) were harvested 1 wk after breaker. For *Aft/Aft atv/atv* fruit, 70 d of storage at 17°C were required to observe 100% of the fruit softened, equivalent to the level of softening observed in Ailsa Craig fruits at 42 d (Fig. 1a,c) and the proportion of fresh weight loss was higher in Ailsa Craig than in *Aft/Aft atv/atv* fruit (Fig. 1b). We repeated the storage test at RT and observed similar results (Fig. 1e, f). After storage for 42 d at RT, the seed in Ailsa Craig fruits showed viviparous germination, followed by complete fruit collapse while *Aft/Aft atv/atv* tomatoes did not (Fig. 1d). The absence of precocious germination was due to elevated anthocyanin levels in the seed of *Aft/Aft atv/atv* plants, rather than elevated levels of

proanthocyanins (Supplementary Information, Fig. S2). The suppression of precocious germination by anthocyanins in the seed has been observed for *Dell/Ros1* purple tomatoes (Butelli *et al.*, 2008) and has been reported following studies of transparent testa mutants in *Arabidopsis* (Abeynayake *et al.*, 2011) and for red wheat compared with white wheat (Flintham, 2000).

Because tomato is a climacteric fruit, ethylene promotes ripening. However, no difference in ethylene production or signaling were detected between high anthocyanin *Dell/Ros1* purple tomatoes and WT tomatoes (Zhang *et al.*, 2013). In addition, due to the light-dependant induction of anthocyanin accumulation of *Aft/Aft atv/atv* fruit, tomatoes grown under natural light have both purple and red skinned regions on the same fruit (Povero *et al.*, 2011). The purple regions have high levels of anthocyanins in the skin, whereas the red regions have very low levels of anthocyanins. The red, low anthocyanin regions underwent normal over-ripening compared with WT Ailsa Craig fruit, and showed more rapid softening than purple regions on the same fruit (Supplementary Information, Fig. S3). This showed that the rate of fruit softening is a localized function associated with anthocyanin production, and therefore not caused by differences in production of the volatile, ethylene. Taken together these results suggest that the accumulation of anthocyanins in the peel of tomato fruits is sufficient to delay post-harvest over-ripening and extend shelf life, although the extension of shelf life was not as great as the doubling observed between purple *Dell/Ros1* tomatoes and their WT controls (Zhang *et al.*, 2013).


Susceptibility to the necrotrophic pathogen *Botrytis cinerea*

The susceptibility of *Aft/Aft atv/atv* fruit to *Botrytis cinerea* was investigated by infecting wounded or intact tomato fruits with fungal spore suspensions. To compare better susceptibility to *B. cinerea* with anthocyanin pigmentation, both purple regions and red regions of fruit grown under natural light were tested. Each *Aft/Aft atv/atv* fruit was sprayed on both purple and red regions or wounded and inoculated with spore cultures of *B. cinerea* strain B05.10. At 3 dpi the proportion of fruits showing symptoms of infection in the purple regions was significantly smaller than for the red regions (Fig. 2a,b). Fungal growth was significantly reduced in *Aft/Aft atv/atv* purple regions compared with growth in the red regions and to growth in the WT line (Ailsa Craig) (Fig. 2c,d). Together, these data demonstrate that resistance is a consequence of anthocyanin accumulation in the purple regions and that anthocyanin pigmentation limited to fruit skin is sufficient to reduce susceptibility to this important necrotrophic pathogen. *Botrytis cinerea* infection induces an oxidative burst by generating reactive oxygen species necessary for pathogen infection (Govrin & Levine, 2000). The reduced susceptibility of anthocyanin-enriched *Aft/Aft atv/atv* fruits could be due to their antioxidant activity, which might counterbalance the oxidative burst induced by the fungus, so limiting pathogen growth (Zhang *et al.*, 2013). Anthocyanin levels are high in *Aft/Aft atv/atv* tomatoes (Mes *et al.*, 2008; Povero *et al.*, 2011) and their presence in anthocyanin-enriched tomato regions correlates with the antioxidant capacity of those regions. Those *Aft/Aft atv/atv* purple fruits that accumulated the highest concentrations of anthocyanins, as a



**Fig. 1** Accumulation of anthocyanins in *Aft/Aft atv/atv* tomatoes (*Solanum lycopersicum*) delays late ripening. Ailsa Craig red (open circles), and *Aft/Aft atv/atv*, purple (closed circles) tomato fruits were stored at 17°C (a–c) and at room temperature (d–f). At 42 d of storage the wild type (WT) fruit showed severe over-ripening symptoms while the *Aft/Aft atv/atv* fruit were still firm (a, d). *Aft/Aft atv/atv* fruits showed slower decrease in fresh weight (FW) compared with red, Ailsa Craig tomatoes (b, e) and slower over-ripening as determined by the percentage of firm fruit (c, f). Fruits were harvested at 7 d post-breaker (d0 = 7 dpb). Fresh weight reduction is presented using the percentage of the initial weight. Error bars, ± standard error of the mean (SEM) (*n* ≥ 8). Percentages of fruit showing over ripening symptoms (softening and shriveling) were assessed visually every week during storage tests.






**Fig. 2** Accumulation of anthocyanins in *Aft/Aft atv/atv* tomatoes (*Solanum lycopersicum*) reduces susceptibility to *Botrytis cinerea*. (a, b) Symptoms of either wounded or sprayed purple and red regions of *Aft/Aft atv/atv* tomatoes fruits after inoculation with *B. cinerea* B05.10. (c) Quantitative PCR revealed more *B. cinerea* growing on the red regions of *Aft/Aft atv/atv* fruits than on purple regions at 3 d post-inoculation (dpi). *B. cinerea* growth was calculated by comparing the ratio of *B. cinerea* DNA to tomato DNA. Error bars, + standard error of the mean (SEM) ( $n = 3$ ).  $*P < 0.05$ , compared with control red regions. (d) The ripening-related increase in susceptibility to *B. cinerea* did not occur in *Aft/Aft atv/atv* purple regions. Lesion diameter was measured 3 dpi. Error bars, + SEM ( $n \geq 3$ ).  $*P < 0.05$ ;  $**P < 0.01$ , for values for purple regions compared with red regions of *Aft/Aft atv/atv* fruits grown under natural light at the same stage of ripening. Ailsa Craig, which does not synthesize anthocyanins in its fruit, was used as control for *B. cinerea* infection.

result of greater exposure to light, had the highest antioxidant capacities (Fig. 3a). Increased cuticle thickness has been reported to be associated with longer shelf life (Yeats *et al.*, 2012), but we observed no significant differences in this trait between *Aft/Aft atv/atv* and Ailsa Craig tomatoes (Fig. 3b). These data suggest that the reduced susceptibility to *B. cinerea* in anthocyanin-enriched fruit is due to their antioxidant content rather than to differences in cuticle thickness.

Accumulation of anthocyanins in tomato fruit skin by genetic modification can extend shelf life

To confirm that the enhanced pathogen resistance observed in *Aft/Aft atv/atv* fruit was due to anthocyanin accumulation and not to another, unknown, trait linked to or associated with either *Aft* or *atv*, we generated tomato lines which accumulated anthocyanins predominantly in skin by genetic modification. Because the *Aft* gene is induced by light, anthocyanins accumulate predominantly in the skin of *Aft/Aft atv/atv* fruit (Supplementary Information, Fig. S4A) (Jones *et al.*, 2003; Mes *et al.*, 2008; Povero *et al.*, 2011).




**Fig. 3** Delayed over-ripening and reduced pathogen susceptibility are associated with the increased antioxidant capacity due to increased anthocyanin levels in *Aft/Aft atv/atv* tomatoes (*Solanum lycopersicum*). (a) Trolox equivalent total antioxidant capacity (TEAC) of water (black bars) and acetone (gray bars) extracts from purple, medium and red regions of *Aft/Aft atv/atv* tomatoes during ripening. Error bars show the standard error of the mean (SEM) ( $n = 3$ ).  $*P < 0.05$ , values for purple regions compared with red regions at the same stage. (b) Cuticle thickness of purple (*Aft/Aft atv/atv*) and red (Ailsa Craig) tomatoes. Measurements were made above the center of each epidermal cell. Error bars, + SEM ( $n \geq 3$ ).

We used the promoter of the *PLI* gene, which is induced by light and is active mainly in tomato skin (Estornell *et al.*, 2009), to drive the expression of the MYB transcriptional factor *Rosea 1*. (Martin *et al.*, 2012) We expressed *PLI:Ros1* together with *35S:Del* in tomato using a binary vector that carried both gene constructs (Martin *et al.*, 2012). The new *PLI:Ros1/35S:Del* (PRD) lines accumulated high levels of anthocyanins in fruit skin, with much less anthocyanin in the fruit flesh compared with previously reported *E8:Del/Ros1* lines (Butelli *et al.*, 2008). From among > 40 independent transgenic lines, two lines, *PRD8-2* and *PRD17-2*, which differentially accumulated anthocyanin in the fruit skin, were selected (Fig. 4a,b). Although both lines accumulated low levels of anthocyanins in flesh, the anthocyanin contents of the flesh of the PRD lines were much lower than in *E8:Del/Ros1* lines (Fig. 4a,b, Supplementary Information, Fig. S4b) When fruit were inoculated with *Botrytis cinerea* culture, both *E8:Del/Ros1* and PRD lines showed smaller lesion size at 3 dpi compared with WT fruits (Fig. 4c). Similar results were observed by spraying intact fruit with *B. cinerea* spores. The proportion of fruit showing severe infection was always lower for the transgenic lines compared with WT fruits (Fig. 4d). In both cases, susceptibility was inversely correlated with anthocyanin content; *E8:Del/Ros1* N and *PRD8-2* tomatoes, which had the highest concentration of anthocyanins, were less susceptible to *B. cinerea* than *PRD17-2* and other transgenic lines. These results showed that accumulation of anthocyanins in tomato skin is sufficient to reduce the susceptibility of fruit to *B. cinerea*.

## Discussion

One of the biggest challenges for the tomato industry is to reduce post-harvest losses resulting from fruit softening and post-harvest infection by several pathogens. So far, biotechnological strategies have been adopted to extend the shelf-life of tomatoes, often at the expense of flavor, aroma, and texture (Baldwin *et al.*, 2011). Anthocyanins, induced by gamma irradiation, have been suggested to prolong the shelf life of grape pomace (Ayed *et al.*, 1999).



**Fig. 4** PRD tomatoes show reduced *Botrytis cinerea* susceptibility. (a) Pictures of different anthocyanin enriched lines: *E8:Del/Ros1 N* and *C*, PRD 8-2 and 17-2 tomatoes (*Solanum lycopersicum*) were taken at the red stage and whole fruit, peeled fruit and skin are shown compared with the wild type Microtom (MT). (b) Anthocyanin levels for all the transgenic tomato lines, error bars,  $\pm$  standard error of the mean (SEM) ( $n = 3$ ). (c) All transgenic lines showed less susceptibility to *B. cinerea* in wound infection tests. Lesion diameter was measured 3 d post-inoculation (dpi). Error bars,  $\pm$  SEM ( $n \geq 3$ ). \* $P < 0.05$ ; \*\* $P < 0.01$ , for values of anthocyanin-enriched tomatoes compared with red Microtom fruit at the same stage. (d) In spray tests the proportion of susceptible tomatoes was low in the anthocyanin-enriched lines, in particular in *E8:Del/Ros1 N* and PRD 8-2, and was inversely correlated with anthocyanin concentration.

Recently, we have shown that genetically modified tomatoes accumulating high levels of anthocyanins in fruit have an extended shelf life compared with controls (Zhang *et al.*, 2013). Here, we show that *Afi/Afi atv/atv* tomato fruit accumulating anthocyanins in the skin have an extended shelf life compared with WT tomatoes. The anthocyanin-enriched sectors of *Afi/Afi atv/atv* tomatoes are less susceptible to *Botrytis cinerea* infection in both wound and spray tests (Fig. 2a,b) and this is correlated to the higher antioxidant capacity of purple tomatoes compared with WT (Fig. 3a). Furthermore, the *Afi/Afi atv/atv* tomatoes showed delayed over-ripening (Fig. 1a,b). Fifty percent softening of *Afi/Afi atv/atv* fruits occurred between 1 and 2 wk later than for WT (Ailsa Craig) tomatoes (Fig. 1c,d) demonstrating an extended shelf life both at 17°C and at RT. Additionally, susceptibility to infection by opportunistic pathogens during storage was higher for red fruit than for purple ones, seeds from *Afi/Afi atv/atv* fruit are not viviparous compared with seed from WT fruit (due to anthocyanin accumulation in *Afi/Afi atv/atv* seed), and *Afi/Afi atv/atv* tomatoes can be stored longer at RT which could reduce the cost of shipping and storage. Taken together, these data show that *Afi/Afi atv/atv* tomatoes have enhanced shelf life due to delayed over-ripening and reduced susceptibility to *B. cinerea*. The increase in shelf life correlated with the presence of anthocyanins and the

antioxidant activity of these anthocyanins could also explain the lower susceptibility to *B. cinerea* (Fig. 3b) (Zhang *et al.*, 2013). To confirm that skin-specific accumulation of anthocyanins in tomato is sufficient to reduce the susceptibility to *B. cinerea* and extend shelf life, we also produced tomatoes which accumulated anthocyanins predominantly in their skin (*PDR* lines). *PDR* fruits, either sprayed or wounded, showed a reduced susceptibility to *B. cinerea* infection (Fig. 4) and susceptibility was inversely correlated with anthocyanin levels. These data strongly support our observations of the extended shelf life of *Afi/Afi atv/atv* tomatoes. This study demonstrates clearly that anthocyanin accumulation in skin is sufficient to reduce susceptibility to *B. cinerea* in tomato fruit. The ability to synthesize anthocyanins in the fruit skin in *Afi/Afi atv/atv* tomatoes could be exploited by breeders to obtain new tomato varieties with both extended shelf life and reduced susceptibility to *B. cinerea*.

### Acknowledgements

L.B. and Y.Z. contributed equally to this work. Y.Z. is supported by a Rotation PhD studentship from the John Innes Foundation. C.M. was supported by the core strategic grant of the Biological and Biotechnological Science Research Council (BBSRC) to the John

Innes Center, and C.M. and Y.Z. are currently supported by the Institute Strategic Program *Understanding and Exploiting Plant and Microbial Secondary Metabolism* (BB/J004596/1) from the BBSRC. H-j.S. was supported by grant BB/G042960/1 from BBSRC and the John Innes Foundation. This work benefited from the networking activities within the European-funded COST ACTION FA1106 QualityFruit.AUTHOR: Please check funding agency and number.

## References

- Abeynayake SW, Panter S, Mouradov A, Spangenberg G. 2011. A high-resolution method for the localization of proanthocyanidins in plant tissues. *Plant Methods* 7: 13.
- Al-sane KO, Povero G, Perata P. 2011. Anthocyanin tomato mutants: overview and characterization of an anthocyanin-less somaclonal mutant. *Plant Biosystems* 145: 436–444.
- Ayed N, Yu HL, Lacroix M. 1999. Improvement of anthocyanin yield and shelf-life extension of grape pomace by gamma irradiation. *Food Research International* 32: 539–543.
- Baldwin E, Plotto A, Narciso J, Bai J. 2011. Effect of 1-methylcyclopropene on tomato flavour components, shelf life and decay as influenced by harvest maturity and storage temperature. *Journal of the Science of Food and Agriculture* 91: 969–980.
- Brummell D, Harpster M. 2001. Cell wall metabolism in fruit softening and quality and its manipulation in transgenic plants. *Plant Molecular Biology* 47: 311–339.
- Buda GJ, Isaacson T, Matas AJ, Paolillo DJ, Rose JKC. 2009. Three-dimensional imaging of plant cuticle architecture using confocal scanning laser microscopy. *Plant Journal* 60: 378–385.
- Buer CS, Imin N, Djordjevic MA. 2010. Flavonoids: new roles for old molecules. *Journal of Integrative Plant Biology* 52: 98–111.
- Butelli E, Titta L, Giorgio M, Mock HP, Matros A, Peterek S, Schijlen EG, Hall RD, Bovy AG, Luo J *et al.* 2008. Enrichment of tomato fruit with health-promoting anthocyanins by expression of select transcription factors. *Nature Biotechnology* 26: 1301–1308.
- De Pascual-Teresa S, Moreno DA, García-Viguera C. 2010. Flavanols and anthocyanins in cardiovascular health: a review of current evidence. *International Journal of Molecular Sciences* 11: 1679–1703.
- Dean R, Van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ, Dickman M, Kahmann R, Ellis J *et al.* 2012. The Top 10 fungal pathogens in molecular plant pathology. *Molecular Plant Pathology* 13: 414–430.
- Estornell LH, Orzáez D, López-Peña L, Pineda B, Antón M, Moreno V, Granell A. 2009. A multisite gateway-based toolkit for targeted gene expression and hairpin RNA silencing in tomato fruits. *Plant Biotechnology Journal* 7: 298–309.
- Fillatti JJ, Kiser J, Rose R, Comai L. 1987. Efficient transfer of a glyphosate tolerance gene into tomato using a binary *Agrobacterium tumefaciens* vector. *Nature Biotechnology* 5: 726–730.
- Flintham J. 2000. Different genetic components control coat-imposed and embryo-imposed dormancy in wheat. *Seed Science Research* 10: 43–50.
- Gonzali S, Mazzucato A, Perata P. 2009. Purple as a tomato: towards high anthocyanin tomatoes. *Trends in Plant Science* 14: 237–241.
- Govrin EM, Levine A. 2000. The hypersensitive response facilitates plant infection by the necrotrophic pathogen *Botrytis cinerea*. *Current Biology* 10: 751–757.
- Grotewold E. 2006. The genetics and biochemistry of floral pigments. *Annual Review of Plant Biology* 57: 761–780.
- Jones CM, Mes P, Myers JR. 2003. Characterization and inheritance of the anthocyanin fruit (Aft) tomato. *Journal of Heredity* 94: 449–456.
- Martin C, Zhang Y, Tomlinson L, Kallam K, Luo J, Jones JDG, Granell A, Orzaez D, Butelli E. 2012. Colouring up plant biotechnology. *Recent Advances in Polyphenol Research* 3: 131–142.
- Mes PJ, Bochens P, Myers JR, Durst R. 2008. Characterization of tomatoes expressing anthocyanin in the fruit. *Journal of the American Society for Horticultural Science* 133: 262–269.
- Nambeesan S, Datsenka T, Ferruzzi MG, Malladi A, Mattoo AK, Handa AK. 2010. Overexpression of yeast spermidine synthase impacts ripening, senescence and decay symptoms in tomato. *Plant Journal* 63: 836–847.
- Pellegrini N, Del Rio D, Colombi B, Bianchi M, Brighenti F. 2003. Application of the 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) radical cation assay to a flow injection system for the evaluation of antioxidant activity of some pure compounds and beverages. *Journal of Agriculture and Food Chemistry* 51: 260–264.
- Povero G, Gonzali S, Bassolino L, Mazzucato A, Perata P. 2011. Transcriptional analysis in high-anthocyanin tomatoes reveals synergistic effect of *Aft* and *atv* genes. *Journal of Plant Physiology* 168: 270–279.
- Sapir M, Oren-Shamir M, Ovadia R, Reuveni M, Evenor D, Tadmor Y, Nahon S, Shlomo H, Chen L, Meir A *et al.* 2008. Molecular aspects of Anthocyanin fruit tomato in relation to high pigment-1. *Journal of Heredity* 99: 292–303.
- Spencer JP. 2010. The impact of fruit flavonoids on memory and cognition. *British Journal of Nutrition* 104(Suppl 3): S40–S47.
- Stefanato FL, Abou-Mansour E, Buchala A, Kretschmer M, Mosbach A, Hahn M, Bochet CG, Metraux JP, Schoonbeek HJ. 2009. The ABC transporter BcarrB from *Botrytis cinerea* exports camalexin and is a virulence factor on *Arabidopsis thaliana*. *Plant Journal* 58: 499–510.
- Valero D, Martinez-Romero DMR, Serrano MS. 2002. The role of polyamines in the improvement of the shelf life of fruit. *Trends in Food Science and Technology* 13: 228–234.
- Vicente AR, Saladie M, Rose JKC, Labavitch JM. 2007. The linkage between cell wall metabolism and fruit softening: looking to the future. *Journal of the Science of Food and Agriculture* 87: 1435–1448.
- Williamson B, Tudzynski B, Tudzynski P, Van Kan JAL. 2007. *Botrytis cinerea*: the cause of grey mould disease. *Molecular Plant Pathology* 8: 561–580.
- Yeats TH, Buda GJ, Wang Z, Chehanovsky N, Moyle LC, Jetter R, Schaffer AA, Rose JK. 2012. The fruit cuticles of wild tomato species exhibit architectural and chemical diversity, providing a new model for studying the evolution of cuticle function. *Plant Journal* 69: 655–666.
- Zhang Y, Butelli E, De Stefano R, Schoonbeek H-J, Magusin A, Pagliarini C, Wellner N, Hill L, Orzaez D, Granell A *et al.* 2013. Anthocyanins double the shelf life of tomatoes by delaying overripening and reducing susceptibility to gray mold. *Current Biology* 23: 1094–1100.

## Supporting Information

Additional supporting information may be found in the online version of this article.

**Fig. S1** High light induces anthocyanin accumulation in the skin of *Aft/Aft atv/atv* fruit.

**Fig. S2** Seeds of *Aft/Aft atv/atv* fruit accumulate anthocyanins.

**Fig. S3** Delayed over-ripening is directly associated with anthocyanin production.

**Fig. S4** Both *Aft/Aft atv/atv* and *PRD* fruit predominantly accumulate anthocyanins in the skin.

Please note: Wiley Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.